

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07K 14/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 99/58558</b> <b>(43) International Publication Date:</b> 18 November 1999 (18.11.99)
<b>(21) International Application Number:</b> PCT/US99/10567 <b>(22) International Filing Date:</b> 13 May 1999 (13.05.99)  <b>(30) Priority Data:</b> 60/085,343 13 May 1998 (13.05.98) US 60/098,010 26 August 1998 (26.08.98) US  <b>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications</b> US 60/085,343 (CIP) Filed on 13 May 1998 (13.05.98) US 60/098,010 (CIP) Filed on 26 August 1998 (26.08.98)  <b>(71) Applicant (for all designated States except US):</b> INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> BANDMAN, Olga [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). HILLMAN, Jennifer, L. [US/US]; 230 Monroe Drive #12, Mountain View, CA 94040 (US). LAL, Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US).		<p>YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). TANG, Y., Tom [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). PATTERSON, Chandra [US/US]; 490 Sherwood Way #1, Menlo Park, CA 94025 (US). BAUGHN, Mariah, R. [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). YANG, Junming [CN/US]; 7136 Clarendon Street, San Jose, CA 95129 (US).</p> <p><b>(74) Agents:</b> BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US).</p> <p><b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i></p>
<b>(54) Title:</b> CELL SIGNALING PROTEINS		
<b>(57) Abstract</b>		
<p>The invention provides human cell signaling proteins (CSIGP) and polynucleotides which identify and encode CSIGP. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating or prevention disorders associated with expression of CSIGP.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## CELL SIGNALING PROTEINS

### TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of cell signaling proteins  
5 and to the use of these sequences in the diagnosis, treatment, and prevention of cell proliferative  
and inflammatory disorders.

### BACKGROUND OF THE INVENTION

10 Signal transduction is the process of biochemical events by which cells respond to  
extracellular signals. Extracellular signals are transduced through a biochemical cascade that  
begins with the binding of a signal molecule such as a hormone, neurotransmitter, or growth  
factor, to a cell membrane receptor and ends with the activation of an intracellular target molecule.  
The process of signal transduction regulates a wide variety of cell functions including cell  
15 proliferation, differentiation, and gene transcription.

Signal transduction is the general process by which cells respond to extracellular signals  
(hormones, neurotransmitters, growth and differentiation factors, etc.) through a cascade of  
biochemical reactions that begins with the binding of the signaling molecule to a cell membrane  
receptor and ends with the activation of an intracellular target molecule. Intermediate steps in this  
20 process involve the activation of various cytoplasmic proteins by phosphorylation via protein  
kinases and the eventual translocation of some of these activated proteins to the cell nucleus where  
the transcription of specific genes is triggered. Thus, the signal transduction process regulates all  
types of cell functions including cell proliferation, differentiation, and gene transcription.

Protein kinases play a key role in the signal transduction process by phosphorylating and  
25 activating various proteins involved in signaling pathways. The high energy phosphate which  
drives this activation is generally transferred from adenosine triphosphate molecules (ATP) to a  
particular protein by protein kinases and removed from that protein by protein phosphatases.  
Phosphorylation occurs in response to extracellular signals, cell cycle checkpoints, and  
environmental or nutritional stresses. Protein kinases are roughly divided into two groups; those  
30 that phosphorylate tyrosine residues (protein tyrosine kinases, PTK) and those that phosphorylate  
serine or threonine residues (serine/threonine kinases, STK). A few protein kinases have dual  
specificity for serine/threonine and tyrosine residues. Almost all kinases contain a similar 250-300  
amino acid catalytic domain containing specific residues and sequence motifs characteristic of the  
kinase family. (Hardie, G. and Hanks, S. (1995) The Protein Kinase Facts Books, Vol I:7-20

Academic Press, San Diego, CA.)

STKs include the second messenger dependent protein kinases such as the cyclic-AMP dependent protein kinases (PKA), which are involved in mediating hormone-induced cellular responses; calcium-calmodulin (CaM) dependent protein kinases, which are involved in regulation of smooth muscle contraction, glycogen breakdown, and neurotransmission; and the mitogen-activated protein kinases (MAP) which mediate signal transduction from the cell surface to the nucleus via phosphorylation cascades. Altered PKA expression is implicated in a variety of disorders and diseases including cancer, thyroid disorders, diabetes, atherosclerosis, and cardiovascular disease. (Isselbacher, K.J. et al. (1994) Harrison's Principles of Internal Medicine, McGraw-Hill, New York, NY, pp. 416-431, 1887.)

PTKs are divided into transmembrane, receptor PTKs and nontransmembrane, non-receptor PTKs. Transmembrane protein-tyrosine kinases are receptors for most growth factors which include epidermal GF, platelet-derived GF, fibroblast GF, hepatocyte GF, insulin and insulin-like GFs, nerve GF, vascular endothelial GF, and macrophage colony stimulating factor. Non-receptor PTKs lack transmembrane regions and, instead, form complexes with the intracellular regions of cell surface receptors. Receptors that function through non-receptor PTKs include those for cytokines, hormones (growth hormone and prolactin) and antigen-specific receptors on T and B lymphocytes.

Many of these PTKs were first identified as the products of mutant oncogenes in cancer cells where their activation was no longer subject to normal cellular controls. In fact, about one third of the known oncogenes encode PTKs, and it is well known that cellular transformation (oncogenesis) is often accompanied by increased tyrosine phosphorylation activity. (Charbonneau H and Tonks NK (1992) *Annu Rev Cell Biol* 8:463-493.)

Protein phosphatases regulate the effects of protein kinases by removing phosphate groups from molecules previously activated by kinases. The two principle categories of protein phosphatases are the protein phosphatases (PPs) and the protein tyrosine phosphatases (PTPs). PPs dephosphorylate phosphoserine/threonine residues and are important regulators of many cAMP-mediated hormone responses in cells. (Cohen, P. (1989) *Annu. Rev. Biochem.* 58:453-508.) PTPs reverse the effects of protein tyrosine kinases and play a significant role in cell cycle and cell signaling processes. (Charbonneau and Tonks, supra.) In the process of cell division, for example, a specific PTP (M-phase inducer phosphatase) plays a key role in the induction of mitosis by dephosphorylating and activating a specific PTK (CDC2) leading to cell division. (Sadu, K.. et al. (1990) *Proc. Natl. Acad. Sci.* 87:5139-5143.)

Guanine nucleotide binding proteins (GTP-binding proteins) are critical mediators of the signal transduction pathway. Extracellular ligands such as hormones, growth factors,

neuromodulators, or other signaling molecules bind to transmembrane receptors, and the signal is propagated to effector molecules by intracellular signal transducing proteins. Many of these signal transduction proteins are GTP-binding proteins which regulate intracellular signaling pathways. GTP-binding proteins participate in a wide range of other regulatory functions including

5 metabolism, growth, differentiation, cytoskeletal organization, and intracellular vesicle transport and secretion. Exchange of bound GDP for GTP followed by hydrolysis of GTP to GDP provides the energy that enables GTP-binding proteins to alter their conformation and interact with other cellular components. Two structurally distinct classes of GTP-binding proteins are recognized: heterotrimeric GTP-binding proteins, consisting of three different subunits, and monomeric, low

10 molecular weight (LMW), GTP-binding proteins consisting of a single polypeptide chain.

G protein coupled receptors (GPCR) are a superfamily of integral membrane proteins which transduce extracellular signals. GPCRs include receptors for biogenic amines, mediators of inflammation, peptide hormones, and sensory signal mediators. A GPCR becomes activated when the receptor binds to its extracellular ligand. The beta subunit of the GPCR, which consists of an

15 amino-terminal helical segment followed by seven WD, or  $\beta$  transducin repeats, transduces signals across the plasma membrane. Conformational changes in the GPCR, resulting from the ligand-receptor interaction, promote the binding of GTP to the GPCR intracellular domains. GTP binding to the GPCR leads to the interaction of the GPCR alpha subunit with adenylate cyclase or other second messenger molecule generators. This interaction regulates the activity of second

20 messenger molecules such as cAMP, cGMP, or eicosinoids which, in turn, regulate phosphorylation and activation of other intracellular proteins. The GPCR changes conformation upon hydrolysis of the bound GTP by GTPases, dissociates from the second messenger molecule generator, and returns to its initial pre-ligand binding conformation.

G beta proteins, also known as  $\beta$  transducins, contain seven tandem repeats of the WD-

25 repeat sequence motif, a motif found in many proteins with regulatory functions. WD-repeat proteins contain from four to eight copies of a loosely conserved repeat of approximately 40 amino acids which participates in protein-protein interactions. Mutations and variant expression of  $\beta$  transducin proteins are linked with various disorders. Mutations in LIS1, a subunit of the human platelet activating factor acetylhydrolase, cause Miller-Dieker lissencephaly. RACK1

30 binds activated protein kinase C, and RbAp48 binds retinoblastoma protein. CstF is required for polyadenylation of mammalian pre-mRNA in vitro and associates with subunits of cleavage-stimulating factor. CD4, an integral membrane glycoprotein which functions as an HIV co-receptor for infection of human host cells is degraded by HIV-encoded Vpu in the endoplasmic reticulum. WD repeats of human beta TrCP molecule mediate the formation of the CD4- Vpu,

35 inducing CD4 proteolysis (Neer, E.J. et al. (1994) Nature 371:297-300 and Margottin, F. et al.

(1998) *Mol. Cell.* 1:565-574).

Irregularities in the GPCR signaling cascade may result in abnormal activation of leukocytes and lymphocytes, leading to the tissue damage and destruction seen in many inflammatory and autoimmune diseases such as rheumatoid arthritis, biliary cirrhosis, hemolytic anemia, lupus erythematosus, and thyroiditis. Abnormal cell proliferation, including cyclic AMP stimulation of brain, thyroid, adrenal, and gonadal tissue proliferation is regulated by G proteins. Mutations in  $G_\alpha$  subunits have been found in growth-hormone-secreting pituitary somatotroph tumors, hyperfunctioning thyroid adenomas, and ovarian and adrenal neoplasms (Meij, J.T.A. (1996) *Mol. Cell. Biochem.* 157:31-38; Aussel, C. et al. (1988) *J. Immunol.* 140:215-220).

10 LMW GTP-binding proteins regulate cell growth, cell cycle control, protein secretion, and intracellular vesicle interaction. They consist of single polypeptides which, like the alpha subunit of the heterotrimeric GTP-binding proteins, are able to bind to and hydrolyze GTP, thus cycling between an inactive and an active state. LMW GTP-binding proteins respond to extracellular signals from receptors and activating proteins by transducing mitogenic signals involved in  
15 various cell functions. The binding and hydrolysis of GTP regulates the response of LMW GTP-binding proteins and acts as an energy source during this process (Bokoch, G. M. and Der, C. J. (1993) *FASEB J.* 7:750-759).

At least sixty members of the LMW GTP-binding protein superfamily have been identified and are currently grouped into the four subfamilies of ras, rho, arf, sar1, ran, and rab.  
20 Activated ras genes were initially found in human cancers and subsequent studies confirmed that ras function is critical in determining whether cells continue to grow or become differentiated. Other members of the LMW G-protein superfamily have roles in signal transduction that vary with the function of the activated genes and the locations of the GTP-binding proteins that initiate the activity. Rho GTP-binding proteins control signal transduction pathways that link growth factor  
25 receptors to actin polymerization, which is necessary for normal cellular growth and division. The rab, arf, and sar1 families of proteins control the translocation of vesicles to and from membranes for protein localization, protein processing, and secretion. Ran GTP-binding proteins are located in the nucleus of cells and have a key role in nuclear protein import, the control of DNA synthesis, and cell-cycle progression (Hall, A. (1990) *Science* 249:635-640; Barbacid, M. (1987) *Ann. Rev*  
30 *Biochem.* 56:779-827; and Sasaki, T. and Takai, Y. (1998) *Biochem. Biophys. Res. Commun.* 245:641-645).

LMW GTP-binding proteins are GTPases which cycle between a GTP-bound active form and a GDP-bound inactive form. This cycle is regulated by proteins that affect GDP dissociation, GTP association, or the rate of GTP hydrolysis. Proteins affecting GDP association are

represented by guanine nucleotide dissociation inhibitors and guanine nucleotide exchange factors (GEP). The best characterized is the mammalian homologue of the *Drosophila* Son-of-Sevenless protein. Proteins affecting GTP hydrolysis are exemplified by GTPase-activating proteins (GAP). Both GEP and GAP activity may be controlled in response to extracellular stimuli and modulated

5 by accessory proteins such as RalBP1 and POB1. The GDP-bound form is converted to the GTP-bound form through a GDP/GTP exchange reaction facilitated by guanine nucleotide-releasing factors. The GTP-bound form is converted to the GDP-bound form by intrinsic GTPase activity, and the conversion is accelerated by GAP (Ikeda, M. et al. (1998) *J. Biol. Chem.* 273:814-821; Quilliam, L. A. (1995) *Bioessays* 17:395-404.). Mutant Ras-family

10 proteins, which bind but can not hydrolyze GTP, are permanently activated, and cause cell proliferation or cancer, as do GEP that activate LMW GTP-binding proteins (Drivas, G. T. et al. (1990) *Mol. Cell. Biol.* 10:1793-1798; and Whitehead, I. P. et al. (1998) *Mol Cell Biol.* 18:4689-4697.)

The discovery of new cell signaling proteins and the polynucleotides encoding them

15 satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of cell proliferative and inflammatory disorders.

#### SUMMARY OF THE INVENTION

20 The invention features substantially purified polypeptides, cell signaling proteins, referred to collectively as "CSIGP" and individually as CSIGP-1 through CSIGP-13. In one aspect, the invention provides a substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-13, and fragments thereof.

The invention further provides a substantially purified variant having at least 90% amino

25 acid identity to at least one of the amino acid sequences selected from the group consisting of SEQ ID NO:1-13, and fragments thereof. The invention also provides an isolated and purified polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-13, and fragments thereof. The invention also includes an isolated and purified polynucleotide variant having at least 70% polynucleotide sequence identity

30 to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-13, and fragments thereof.

Additionally, the invention provides an isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-13, and fragments

35 thereof. The invention also provides an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide encoding the polypeptide comprising the amino

acid sequence selected from the group consisting of SEQ ID NO:1-13, and fragments thereof.

The invention also provides an isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:14-26, and fragments thereof. The invention further provides an isolated and purified polynucleotide variant having at least 70% polynucleotide sequence identity to the polynucleotide sequence selected from the group consisting of SEQ ID NO:14-26 and fragments thereof. The invention also provides an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:14-26 and fragments thereof.

The invention also provides a method for detecting a polynucleotide in a sample containing nucleic acids, the method comprising the steps of (a) hybridizing the complement of the polynucleotide sequence to at least one of the polynucleotides of the sample, thereby forming a hybridization complex; and (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of a polynucleotide in the sample. In one aspect, the method further comprises amplifying the polynucleotide prior to hybridization.

The invention further provides an expression vector containing at least a fragment of the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-13, and fragments thereof. In another aspect, the expression vector is contained within a host cell.

The invention also provides a method for producing a polypeptide, the method comprising the steps of: (a) culturing the host cell containing an expression vector containing at least a fragment of a polynucleotide under conditions suitable for the expression of the polypeptide; and (b) recovering the polypeptide from the host cell culture.

The invention also provides a pharmaceutical composition comprising a substantially purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1-13, and fragments thereof, in conjunction with a suitable pharmaceutical carrier.

The invention further includes a purified antibody which binds to a polypeptide selected from the group consisting of SEQ ID NO:1-13, and fragments thereof. The invention also provides a purified agonist and a purified antagonist to the polypeptide.

The invention also provides a method for treating or preventing a disorder associated with decreased expression or activity of CSIGP, the method comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising a substantially purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1-13, and fragments thereof, in conjunction with a suitable pharmaceutical carrier.



The invention also provides a method for treating or preventing a disorder associated with increased expression or activity of CSIGP, the method comprising administering to a subject in need of such treatment an effective amount of an antagonist of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-13, and fragments thereof, in  
5 conjunction with a suitable pharmaceutical carrier.

#### BRIEF DESCRIPTION OF THE TABLES

Table 1 shows nucleotide and polypeptide sequence identification numbers (SEQ ID NO), clone identification numbers (clone ID), cDNA libraries, and cDNA fragments used to assemble  
10 full-length sequences encoding CSIGP.

Table 2 shows features of each polypeptide sequence including potential motifs, homologous sequences, and methods and algorithms used for identification of CSIGP.

Table 3 shows the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis, diseases, disorders or conditions associated with these tissues,  
15 and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which Incyte cDNA clones encoding CSIGP were isolated.

Table 5 shows the programs, their descriptions, references, and threshold parameters used to analyze CSIGP.  
20

#### DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the  
25 purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an  
30 antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described  
35 herein can be used to practice or test the present invention, the preferred machines, materials and

methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of  
5 prior invention.

## DEFINITIONS

"CSIGP" refers to the amino acid sequences of substantially purified CSIGP obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and preferably the human species, from any source, whether natural, synthetic,  
10 semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which, when bound to CSIGP, increases or prolongs the duration of the effect of CSIGP. Agonists may include proteins, nucleic acids, carbohydrates, or any other molecules which bind to and modulate the effect of CSIGP.

An "allelic variant" is an alternative form of the gene encoding CSIGP. Allelic variants  
15 may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. Any given natural or recombinant gene may have none, one, or many allelic forms. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination  
20 with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding CSIGP include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polynucleotide the same as CSIGP or a polypeptide with at least one functional characteristic of CSIGP. Included within this definition are polymorphisms which may or may not be readily detectable using a particular  
25 oligonucleotide probe of the polynucleotide encoding CSIGP, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding CSIGP. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent CSIGP. Deliberate amino acid substitutions may be made  
30 on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of CSIGP is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, positively charged amino acids may include lysine and arginine, and amino acids with uncharged polar head groups having similar hydrophilicity values may include leucine, isoleucine,  
35 and valine; glycine and alanine; asparagine and glutamine; serine and threonine; and

phenylalanine and tyrosine.

The terms "amino acid" or "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. In this context, "fragments," "immunogenic fragments," or "antigenic fragments" refer to fragments of CSIGP which are preferably at least 5 to about 15 amino acids in length, most preferably at least 14 amino acids, and which retain some biological activity or immunological activity of CSIGP. Where "amino acid sequence" is recited to refer to an amino acid sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which, when bound to CSIGP, decreases the amount or the duration of the effect of the biological or immunological activity of CSIGP. Antagonists may include proteins, nucleic acids, carbohydrates, antibodies, or any other molecules which decrease the effect of CSIGP.

The term "antibody" refers to intact molecules as well as to fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv fragments, which are capable of binding the epitopic determinant. Antibodies that bind CSIGP polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that fragment of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (given regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition containing a nucleic acid sequence which is complementary to the "sense" strand of a specific nucleic acid sequence. Antisense molecules may be produced by any method including synthesis or transcription. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form

duplexes and to block either transcription or translation. The designation "negative" can refer to the antisense strand, and the designation "positive" can refer to the sense strand.

The term "biologically active," refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic CSIGP, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" or "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence "5' A-G-T 3'" bonds to the complementary sequence "3' T-C-A 5'." Complementarity between two single-stranded molecules may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acids strands, and in the design and use of peptide nucleic acid (PNA) molecules.

A "composition comprising a given polynucleotide sequence" or a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding CSIGP or fragments of CSIGP may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, extended using XL-PCR kit (Perkin-Elmer, Norwalk CT) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of more than one Incyte Clone using a computer program for fragment assembly, such as the GELVIEW Fragment Assembly system (GCG, Madison WI). Some sequences have been both extended and assembled to produce the consensus sequence.

The term "correlates with expression of a polynucleotide" indicates that the detection of the presence of nucleic acids, the same or related to a nucleic acid sequence encoding CSIGP, by northern analysis is indicative of the presence of nucleic acids encoding CSIGP in a sample, and thereby correlates with expression of the transcript from the polynucleotide encoding CSIGP.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to the chemical modification of a polypeptide sequence, or a polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

The term "similarity" refers to a degree of complementarity. There may be partial similarity or complete similarity. The word "identity" may substitute for the word "similarity." A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid is referred to as "substantially similar." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially similar sequence or hybridization probe will compete for and inhibit the binding of a completely similar (identical) sequence to the target sequence under conditions of reduced stringency. This is not to say that conditions of reduced stringency are such that non-specific binding is permitted, as reduced stringency conditions require that the binding of two sequences to one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% similarity or identity). In the absence of non-specific binding, the substantially similar sequence or probe will not hybridize to the second non-complementary target sequence.

The phrases "percent identity" or "% identity" refer to the percentage of sequence similarity found in a comparison of two or more amino acid or nucleic acid sequences. Percent identity can be determined electronically, e.g., by using the MEGALIGN program (DNASTAR, Madison WI). The MEGALIGN program can create alignments between two or more sequences according to different methods, e.g., the clustal method. (See, e.g., Higgins, D.G. and P.M. Sharp (1988) Gene 73:237-244.) The clustal algorithm groups sequences into clusters by examining the distances between all pairs. The clusters are aligned pairwise and then in groups. The percentage similarity between two amino acid sequences, e.g., sequence A and sequence B, is calculated by dividing the length of sequence A, minus the number of gap residues in sequence A, minus the number of gap residues in sequence B, into the sum of the residue matches between sequence A and sequence B, times one hundred. Gaps of low or of no similarity between the two amino acid

sequences are not included in determining percentage similarity. Percent identity between nucleic acid sequences can also be counted or calculated by other methods known in the art, e.g., the Jotun Hein method. (See, e.g., Hein, J. (1990) *Methods Enzymol.* 183:626-645.) Identity between sequences can also be determined by other methods known in the art, e.g., by varying

5 hybridization conditions.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for stable mitotic chromosome segregation and maintenance.

The term "humanized antibody" refers to antibody molecules in which the amino acid  
10 sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to any process by which a strand of nucleic acid binds with a complementary strand through base pairing.

The term "hybridization complex" refers to a complex formed between two nucleic acid  
15 sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g.,  $C_0t$  or  $R_0t$  analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

20 The words "insertion" or "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively, to the sequence found in the naturally occurring molecule.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by  
25 expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

The term "microarray" refers to an arrangement of distinct polynucleotides on a substrate.

The terms "element" or "array element" in a microarray context, refer to hybridizable polynucleotides arranged on the surface of a substrate.

30 The term "modulate" refers to a change in the activity of CSIGP. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of CSIGP.

The phrases "nucleic acid" or "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or  
35 RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may

represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material. In this context, "fragments" refers to those nucleic acid sequences which, when translated, would produce polypeptides retaining some functional characteristic, e.g., antigenicity, or structural domain characteristic, e.g., ATP-binding site, of the full-length  
5 polypeptide.

The terms "operably associated" or "operably linked" refer to functionally related nucleic acid sequences. A promoter is operably associated or operably linked with a coding sequence if the promoter controls the translation of the encoded polypeptide. While operably associated or operably linked nucleic acid sequences can be contiguous and in the same reading frame, certain  
10 genetic elements, e.g., repressor genes, are not contiguously linked to the sequence encoding the polypeptide but still bind to operator sequences that control expression of the polypeptide.

The term "oligonucleotide" refers to a nucleic acid sequence of at least about 6 nucleotides to 60 nucleotides, preferably about 15 to 30 nucleotides, and most preferably about 20 to 25 nucleotides, which can be used in PCR amplification or in a hybridization assay or  
15 microarray. "Oligonucleotide" is substantially equivalent to the terms "amplimer," "primer," "oligomer," and "probe," as these terms are commonly defined in the art.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition.  
20 PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic acids encoding CSIGP, or fragments thereof, or CSIGP itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic  
25 DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" or "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, or an antagonist. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the  
30 presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "stringent conditions" refers to conditions which permit hybridization between polynucleotides and the claimed polynucleotides. Stringent conditions can be defined by salt  
35 concentration, the concentration of organic solvent, e.g., formamide, temperature, and other

conditions well known in the art. In particular, stringency can be increased by reducing the concentration of salt, increasing the concentration of formamide, or raising the hybridization temperature.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least about 60% free, preferably about 75% free, and most preferably about 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

10 "Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

"Transformation" describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle bombardment.

15 20 The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "variant" of CSIGP polypeptides refers to an amino acid sequence that is altered by one or more amino acid residues. The variant may have "conservative" changes, wherein a substituted amino acid has similar structural or chemical properties (e.g., replacement of leucine with isoleucine). More rarely, a variant may have "nonconservative" changes (e.g., replacement of glycine with tryptophan). Analogous minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be substituted, inserted, or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, LASERGENE software (DNASTAR).

25 30

The term "variant," when used in the context of a polynucleotide sequence, may encompass a polynucleotide sequence related to CSIGP. This definition may also include, for example, "allelic" (as defined above), "splice," "species," or "polymorphic" variants. A splice variant may have significant identity to a reference molecule, but will generally have a greater or

35



lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or an absence of domains.

Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A

- 5 polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

## 10 THE INVENTION

The invention is based on the discovery of new human cell signaling proteins (CSIGP), the polynucleotides encoding CSIGP, and the use of these compositions for the diagnosis, treatment, or prevention of cell proliferative and inflammatory disorders.

- Table 1 lists the Incyte Clones used to derive full length nucleotide sequences encoding CSIGP. Columns 1 and 2 show the sequence identification numbers (SEQ ID NO) of the amino acid and nucleic acid sequences, respectively. Column 3 shows the Clone ID of the Incyte Clone in which nucleic acids encoding each CSIGP were first identified, and column 4, the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones, their corresponding cDNA libraries, and shotgun sequences useful as fragments in hybridization technologies, and which are part of the consensus nucleotide sequence of each CSIGP.

- The columns of Table 2 show various properties of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3, potential phosphorylation sites; column 4, potential glycosylation sites; column 5, the amino acid residues comprising signature sequences and motifs; column 6, homologous sequences; and column 7, analytical methods used to identify each protein through sequence homology and protein motifs.

- The columns of Table 3 show the tissue-specificity and disease-association of nucleotide sequences encoding CSIGP. The first column of Table 3 lists the polynucleotide sequence identifiers. The second column lists tissue categories which express CSIGP as a fraction of total tissue categories expressing CSIGP. The third column lists diseases, disorders, and conditions associated with those tissues expressing CSIGP. The fourth column lists the vectors used to subclone the cDNA library.

- The following fragments of the nucleotide sequences encoding CSIGP are useful in hybridization or amplification technologies to identify SEQ ID NO:14-26 and to distinguish between SEQ ID NO:14-26 and similar polynucleotide sequences. The useful fragments are the

fragment of SEQ ID NO:14 from about nucleotide 135 to about nucleotide 189, the  
fragment of SEQ ID NO:15 from about nucleotide 493 to about nucleotide 558, the  
fragment of SEQ ID NO:16 from about nucleotide 1170 to about nucleotide 1233, the  
fragment of SEQ ID NO:17 from about nucleotide 939 to about nucleotide 996, the  
5 fragment of SEQ ID NO:18 from about nucleotide 424 to about nucleotide 486, the  
fragment of SEQ ID NO:19 from about nucleotide 274 to about nucleotide 333, and the  
fragment of SEQ ID NO:20 from about nucleotide 1013 to about nucleotide 1070, the  
fragment of SEQ ID NO:21 from about nucleotide 284 to about nucleotide 325, the fragment of  
SEQ ID NO:22 from about nucleotide 642 to about nucleotide 674, the fragment of SEQ ID  
10 NO:23 from about nucleotide 742 to about nucleotide 769, the fragment of SEQ ID NO:24 from  
about nucleotide 457 to about nucleotide 486, the fragment of SEQ ID NO:25 from about  
nucleotide 205 to about nucleotide 246, and the fragment of SEQ ID NO:26 from about nucleotide  
319 to about nucleotide 342.

The invention also encompasses CSIGP variants. A preferred CSIGP variant is one which  
15 has at least about 80%, more preferably at least about 90%, and most preferably at least about 95%  
amino acid sequence identity to the CSIGP amino acid sequence, and which contains at least one  
functional or structural characteristic of CSIGP.

The invention also encompasses polynucleotides which encode CSIGP. In a particular  
embodiment, the invention encompasses a polynucleotide sequence comprising a sequence  
20 selected from the group consisting of SEQ ID NO:14-26 which encodes CSIGP.

The invention also encompasses a variant of a polynucleotide sequence encoding CSIGP.  
In particular, such a variant polynucleotide sequence will have at least about 70%, more preferably  
at least about 85%, and most preferably at least about 95% polynucleotide sequence identity to the  
polynucleotide sequence encoding CSIGP. A particular aspect of the invention encompasses a  
25 variant of a polynucleotide sequence comprising a sequence selected from the group consisting of  
SEQ ID NO:14-26 which has at least about 70%, more preferably at least about 85%, and most  
preferably at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected  
from the group consisting of SEQ ID NO:14-26. Any one of the polynucleotide variants described  
above can encode an amino acid sequence which contains at least one functional or structural  
30 characteristic of CSIGP

It will be appreciated by those skilled in the art that as a result of the degeneracy of the  
genetic code, a multitude of polynucleotide sequences encoding CSIGP, some bearing minimal  
similarity to the polynucleotide sequences of any known and naturally occurring gene, may be  
produced. Thus, the invention contemplates each and every possible variation of polynucleotide

sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring CSIGP, and all such variations are to be considered as being specifically disclosed.

5           Although nucleotide sequences which encode CSIGP and its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring CSIGP under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding CSIGP or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at  
10   which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding CSIGP and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally  
15   occurring sequence.

          The invention also encompasses production of DNA sequences which encode CSIGP and CSIGP derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to  
20   introduce mutations into a sequence encoding CSIGP or any fragment thereof.

          Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:14-26 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.*  
25   152:507-511.) For example, stringent salt concentration will ordinarily be less than about 750 mM NaCl and 75 mM trisodium citrate, preferably less than about 500 mM NaCl and 50 mM trisodium citrate, and most preferably less than about 250 mM NaCl and 25 mM trisodium citrate. Low stringency hybridization can be obtained in the absence of organic solvent, e.g., formamide, while high stringency hybridization can be obtained in the presence of at least about 35% formamide,  
30   and most preferably at least about 50% formamide. Stringent temperature conditions will ordinarily include temperatures of at least about 30°C, more preferably of at least about 37°C, and most preferably of at least about 42°C. Varying additional parameters, such as hybridization time, the concentration of detergent, e.g., sodium dodecyl sulfate (SDS), and the inclusion or exclusion of carrier DNA, are well known to those skilled in the art. Various levels of stringency are  
35   accomplished by combining these various conditions as needed. In a preferred embodiment,

hybridization will occur at 30°C in 750 mM NaCl, 75 mM trisodium citrate, and 1% SDS. In a more preferred embodiment, hybridization will occur at 37°C in 500 mM NaCl, 50 mM trisodium citrate, 1% SDS, 35% formamide, and 100 µg/ml denatured salmon sperm DNA (ssDNA). In a most preferred embodiment, hybridization will occur at 42°C in 250 mM NaCl, 25 mM trisodium citrate, 1% SDS, 50 % formamide, and 200 µg/ml ssDNA. Useful variations on these conditions will be readily apparent to those skilled in the art.

The washing steps which follow hybridization can also vary in stringency. Wash stringency conditions can be defined by salt concentration and by temperature. As above, wash stringency can be increased by decreasing salt concentration or by increasing temperature. For example, stringent salt concentration for the wash steps will preferably be less than about 30 mM NaCl and 3 mM trisodium citrate, and most preferably less than about 15 mM NaCl and 1.5 mM trisodium citrate. Stringent temperature conditions for the wash steps will ordinarily include temperature of at least about 25°C, more preferably of at least about 42°C, and most preferably of at least about 68°C. In a preferred embodiment, wash steps will occur at 25°C in 30 mM NaCl, 3 mM trisodium citrate, and 0.1% SDS. In a more preferred embodiment, wash steps will occur at 42°C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS. In a most preferred embodiment, wash steps will occur at 68°C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS. Additional variations on these conditions will be readily apparent to those skilled in the art.

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Perkin-Elmer), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the HYDRA microdispenser (Robbins Scientific, Sunnyvale CA), MICROLAB 2200 (Hamilton, Reno NV), Peltier Thermal Cyclers 200 (PTC200; MJ Research, Watertown MA) and the ABI CATALYST 800 (Perkin-Elmer). Sequencing is then carried out using either ABI 373 or 377 DNA Sequencing Systems (Perkin-Elmer) or the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA). The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding CSIGP may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect

upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent  
5 directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In  
10 this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-306). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic  
15 DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

20 When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

25 Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal  
30 using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Perkin-Elmer), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof  
35 which encode CSIGP may be cloned in recombinant DNA molecules that direct expression of

CSIGP, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express CSIGP.

5           The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter CSIGP-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example,  
10 oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

          In another embodiment, sequences encoding CSIGP may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucl.  
15 Acids Res. Symp. Ser. 215-223, and Horn, T. et al. (1980) Nucl. Acids Res. Symp. Ser. 225-232.) Alternatively, CSIGP itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be achieved using the ABI 431A Peptide Synthesizer (Perkin-Elmer). Additionally, the amino acid sequence of  
20 CSIGP, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

          The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) Methods Enzymol. 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by  
25 sequencing. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY.)

          In order to express a biologically active CSIGP, the nucleotide sequences encoding CSIGP or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted  
30 coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding CSIGP. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding CSIGP. Such signals include the ATG initiation codon and adjacent  
35 sequences, e.g. the Kozak sequence. In cases where sequences encoding CSIGP and its initiation

codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous  
5 translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct  
10 expression vectors containing sequences encoding CSIGP and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons,  
15 New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding CSIGP. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral  
20 expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected  
25 depending upon the use intended for polynucleotide sequences encoding CSIGP. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding CSIGP can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or pSPORT1 plasmid (Life Technologies). Ligation of sequences encoding CSIGP into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure  
30 for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509.) When large quantities of CSIGP are needed, e.g. for the production of antibodies, vectors which direct high level expression of CSIGP  
35 may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage

promoter may be used.

Yeast expression systems may be used for production of CSIGP. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors  
5 direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra; Grant et al. (1987) *Methods Enzymol.* 153:516-54; and Scorer, C. A. et al. (1994) *Bio/Technology* 12:181-184.)

Plant systems may also be used for expression of CSIGP. Transcription of sequences  
10 encoding CSIGP may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell*  
15 *Differ.* 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding CSIGP may be ligated  
20 into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses CSIGP in host cells. (See, e.g., Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci.* 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.  
25 SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) *Nat Genet.*  
30 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of CSIGP in cell lines is preferred. For example, sequences encoding CSIGP can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate  
35 vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2



days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

- 5 Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk*<sup>-</sup> or *apr*<sup>-</sup> cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers
- 10 resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* or *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g.,
- 15 Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech),  $\beta$  glucuronidase and its substrate  $\beta$ -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol.
- 20 Biol. 55:121-131.)

- Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding CSIGP is inserted within a marker gene sequence, transformed cells containing sequences encoding CSIGP can be identified by the absence of marker gene
- 25 function. Alternatively, a marker gene can be placed in tandem with a sequence encoding CSIGP under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

- In general, host cells that contain the nucleic acid sequence encoding CSIGP and that express CSIGP may be identified by a variety of procedures known to those of skill in the art.
- 30 These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

- Immunological methods for detecting and measuring the expression of CSIGP using either
- 35 specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques

include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on CSIGP is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art.

- 5 (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St Paul MN, Sect. IV; Coligan, J. E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols. Humana Press, Totowa NJ).

A wide variety of labels and conjugation techniques are known by those skilled in the art  
10 and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding CSIGP include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding CSIGP, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are  
15 commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes,  
20 fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding CSIGP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the  
25 sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode CSIGP may be designed to contain signal sequences which direct secretion of CSIGP through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications  
30 of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38), are available from  
35 the American Type Culture Collection (ATCC, Bethesda MD) and may be chosen to ensure the

correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding CSIGP may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric CSIGP protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of CSIGP activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the CSIGP encoding sequence and the heterologous protein sequence, so that CSIGP may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled CSIGP may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract systems (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, preferably <sup>35</sup>S-methionine.

Fragments of CSIGP may be produced not only by recombinant production, but also by direct peptide synthesis using solid-phase techniques. (See, e.g., Creighton, supra pp. 55-60.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin-Elmer). Various fragments of CSIGP may be synthesized separately and then combined to produce the full length molecule.

## THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between CSIGP and cell signaling proteins. In addition, the expression of CSIGP is closely associated with cell proliferation and inflammatory disorders. Therefore, in cell proliferative and

inflammatory disorders where CSIGP is an inhibitor or suppressor of cell proliferation, it is desirable to increase the expression of CSIGP. In cell proliferative and inflammatory disorders where CSIGP is an activator or enhancer and is promoting cell proliferation, it is desirable to decrease the expression of CSIGP.

5           Therefore, in one embodiment, CSIGP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of CSIGP. Examples of such disorders include, but are not limited to, a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, 10 polycythemia vera, psoriasis, primary thrombocythemia; cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; and an inflammatory 15 disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema 20 nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, 25 thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma.

          In another embodiment, a vector capable of expressing CSIGP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased 30 expression or activity of CSIGP including, but not limited to, those described above.

          In a further embodiment, a pharmaceutical composition comprising a substantially purified CSIGP in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of CSIGP including, but not limited to, those provided above.

35           In still another embodiment, an agonist which modulates the activity of CSIGP may be

administered to a subject to treat or prevent a disorder associated with decreased expression or activity of CSIGP including, but not limited to, those listed above.

In a further embodiment, an antagonist of CSIGP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of CSIGP. Examples of  
5 such disorders include, but are not limited to, those described above. In one aspect, an antibody which specifically binds CSIGP may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissue which express CSIGP.

In an additional embodiment, a vector expressing the complement of the polynucleotide  
10 encoding CSIGP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of CSIGP including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination  
15 therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

20 An antagonist of CSIGP may be produced using methods which are generally known in the art. In particular, purified CSIGP may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind CSIGP. Antibodies to CSIGP may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments,  
25 and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are especially preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with CSIGP or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various  
30 adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

35 It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to

CSIGP have an amino acid sequence consisting of at least about 5 amino acids, and, more preferably, of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of

5 CSIGP amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to CSIGP may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-

10 hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci.* 80:2026-2030; and Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate

15 antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci.* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda, S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce CSIGP-specific single chain antibodies. Antibodies with related specificity, but of distinct

20 idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton D.R. (1991) *Proc. Natl. Acad. Sci.* 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci.* 86:

25 3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299.)

Antibody fragments which contain specific binding sites for CSIGP may also be generated. For example, such fragments include, but are not limited to, F(ab')<sub>2</sub> fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragments. Alternatively, Fab expression libraries may be

30 constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) *Science* 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in

35 the art. Such immunoassays typically involve the measurement of complex formation between

CSIGP and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering CSIGP epitopes is preferred, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for ABBR. Affinity is expressed as an association constant,  $K_a$ , which is defined as the molar concentration of ABBR-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The  $K_a$  determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple ABBR epitopes, represents the average affinity, or avidity, of the antibodies for ABBR. The  $K_a$  determined for a preparation of monoclonal antibodies, which are monospecific for a particular ABBR epitope, represents a true measure of affinity. High-affinity antibody preparations with  $K_a$  ranging from about  $10^9$  to  $10^{12}$  L/mole are preferred for use in immunoassays in which the ABBR-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with  $K_a$  ranging from about  $10^6$  to  $10^7$  L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of ABBR, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J. E. and Cryer, A. (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is preferred for use in procedures requiring precipitation of ABBR-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding CSIGP, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding CSIGP may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding CSIGP. Thus, complementary molecules or fragments may be used to modulate CSIGP activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding CSIGP.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses,

or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences complementary to the polynucleotides encoding CSIGP. (See, e.g., Sambrook, supra; Ausubel, 1995, supra.)

5 Genes encoding CSIGP can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding CSIGP. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a  
10 month or more with a non-replicating vector, and may last even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding CSIGP. Oligonucleotides derived from the transcription  
15 initiation site, e.g., between about positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al.  
20 (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the  
25 ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding CSIGP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences:  
30 GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

35 Complementary ribonucleic acid molecules and ribozymes of the invention may be



prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding CSIGP. Such DNA sequences may be  
5 incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3'  
10 ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and  
15 uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers  
20 may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nature Biotechnology 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

25 An additional embodiment of the invention relates to the administration of a pharmaceutical or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of CSIGP, antibodies to CSIGP, and mimetics, agonists, antagonists, or inhibitors of CSIGP. The compositions may be administered alone or in combination with at least one other agent, such as a  
30 stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial,  
35 intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal,

enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used

5 pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets,  
10 pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combining active compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable  
15 excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and tragacanth; and proteins, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone,  
20 agar, and alginic acid or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for  
25 product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with fillers or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft  
30 capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain  
35 substances which increase the viscosity of the suspension, such as sodium carboxymethyl

cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate, triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the  
5 suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a  
10 manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acid. Salts tend to be more soluble in aqueous or other protonic solvents than are the  
15 corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 mM histidine, 0.1% to 2% sucrose, and 2% to 7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an  
20 appropriate container and labeled for treatment of an indicated condition. For administration of CSIGP, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the  
25 art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes  
30 for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example CSIGP or fragments thereof, antibodies of CSIGP, and agonists, antagonists or inhibitors of CSIGP, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals,  
35 such as by calculating the ED<sub>50</sub> (the dose therapeutically effective in 50% of the population) or

LD<sub>50</sub> (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the LD<sub>50</sub>/ED<sub>50</sub> ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED<sub>50</sub> with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 µg to 100,000 µg, up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

## DIAGNOSTICS

In another embodiment, antibodies which specifically bind CSIGP may be used for the diagnosis of cell proliferative and inflammatory disorders characterized by expression of CSIGP, or in assays to monitor patients being treated with CSIGP or agonists, antagonists, or inhibitors of CSIGP. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for CSIGP include methods which utilize the antibody and a label to detect CSIGP in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring CSIGP, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of CSIGP expression. Normal or standard values for CSIGP expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to

CSIGP under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, preferably by photometric means. Quantities of CSIGP expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for  
5 diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding CSIGP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantitate gene expression in biopsied tissues in which expression of CSIGP  
10 may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of CSIGP, and to monitor regulation of CSIGP levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding CSIGP or closely related  
15 molecules may be used to identify nucleic acid sequences which encode CSIGP. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification (maximal, high, intermediate, or low), will determine whether the probe identifies only naturally occurring sequences encoding CSIGP, allelic variants, or related sequences.

20 Probes may also be used for the detection of related sequences, and should preferably have at least 50% sequence identity to any of the CSIGP encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:14-26 or from genomic sequences including promoters, enhancers, and introns of the CSIGP gene.

25 Means for producing specific hybridization probes for DNAs encoding CSIGP include the cloning of polynucleotide sequences encoding CSIGP or CSIGP derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a  
30 variety of reporter groups, for example, by radionuclides such as <sup>32</sup>P or <sup>35</sup>S, or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding CSIGP may be used for the diagnosis of cell proliferative and inflammatory disorders associated with expression of CSIGP. Examples of such disorders include, but are not limited to, a disorder of cell proliferation such as actinic keratosis,  
35 arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease

(MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia; cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; and an inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma. The polynucleotide sequences encoding CSIGP may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and ELISA assays; and in microarrays utilizing fluids or tissues from patients to detect altered CSIGP expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding CSIGP may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding CSIGP may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding CSIGP in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of CSIGP, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a

sequence, or a fragment thereof, encoding CSIGP, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with  
5 values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results  
10 obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A  
15 more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding CSIGP may involve the use of PCR. These oligomers may be chemically synthesized, generated  
20 enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding CSIGP, or a fragment of a polynucleotide complementary to the polynucleotide encoding CSIGP, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantitation of closely related DNA or RNA sequences.

25 Methods which may also be used to quantitate the expression of CSIGP include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in an ELISA format  
30 where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously  
35 and to identify genetic variants, mutations, and polymorphisms. This information may be used to

determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)

In another embodiment of the invention, nucleic acid sequences encoding CSIGP may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) site. Correlation between the location of the gene encoding CSIGP on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.



In another embodiment of the invention, CSIGP, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of  
5 binding complexes between CSIGP and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with CSIGP, or  
10 fragments thereof, and washed. Bound CSIGP is then detected by methods well known in the art. Purified CSIGP can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which  
15 neutralizing antibodies capable of binding CSIGP specifically compete with a test compound for binding CSIGP. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with CSIGP.

In additional embodiments, the nucleotide sequences which encode CSIGP may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely  
20 on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of  
25 the remainder of the disclosure in any way whatsoever.

The entire disclosure of all applications, patents, and publications, cited above and below, and of US provisional applications 60/085,343 (filed May 13, 1998), and 60/098,010 (filed August 26, 1998) are hereby incorporated by reference.

## EXAMPLES

### 30 I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting  
35 lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated

from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A<sup>+</sup>) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Valencia CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, *supra*, units 5.1-6.6). Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), pSPORT1 plasmid (Life Technologies), or pINCY (Incyte Pharmaceuticals, Palo Alto CA). Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 $\alpha$ , DH10B, or ElectroMAX DH10B from Life Technologies.

## II. Isolation of cDNA Clones

Plasmids were recovered from host cells by *in vivo* excision, using the UNIZAP vector system (Stratagene) or cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the REAL Prep 96 plasmid kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a Fluoroskan II

fluorescence scanner (Labsystems Oy, Helsinki, Finland).

### III. Sequencing and Analysis

The cDNAs were prepared for sequencing using either an ABI CATALYST 800 (Perkin-Elmer) or a HYDRA microdispenser (Robbins) or MICROLAB 2200 (Hamilton) sequencing  
5 preparation system in combination with PTC-200 thermal cyclers (MJ Research). The cDNAs were sequenced using the ABI PRISM 373 or 377 sequencing systems of the MEGABACE 1000 DNA sequencing system (Molecular Dynamics) and ABI protocols, base calling software, and kits (Perkin-Elmer). Alternatively, solutions and dyes from Amersham Pharmacia Biotech were used. Reading frames were determined using standard methods (Ausubel, 1997, supra). Some of the  
10 cDNA sequences were selected for extension using the techniques disclosed in Example V.

The polynucleotide sequences derived from cDNA, extension, and shotgun sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the software programs, descriptions, references, and threshold parameters used. The first column of Table 5 shows the tools, programs,  
15 and algorithms used, the second column provides a brief description thereof, the third column presents the references which are incorporated by reference herein, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the probability the greater the homology). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software  
20 Engineering, S. San Francisco CA) and LASERGENE software (DNASTAR).

cDNAs were also compared to sequences in GenBank using a search algorithm developed by Applied Biosystems and incorporated into the INHERIT™ 670 sequence analysis system. In this algorithm, Pattern Specification Language (TRW Inc, Los Angeles, CA) was used to determine regions of homology. The three parameters that determine how the sequence  
25 comparisons run were window size, window offset, and error tolerance. Using a combination of these three parameters, the DNA database was searched for sequences containing regions of homology to the query sequence, and the appropriate sequences were scored with an initial value. Subsequently, these homologous regions were examined using dot matrix homology plots to distinguish regions of homology from chance matches. Smith-Waterman alignments were used to  
30 display the results of the homology search.

Peptide and protein sequence homologies were ascertained using the INHERIT- 670 sequence analysis system using the methods similar to those used in DNA sequence homologies. Pattern Specification Language and parameter windows were used to search protein databases for sequences containing regions of homology which were scored with an initial value. Dot-matrix  
35 homology plots were examined to distinguish regions of significant homology from chance

matches.

The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programing, and dinucleotide nearest neighbor analysis. The sequences were then  
 5 queried against a selection of public databases such as GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS to acquire annotation, using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length  
 10 polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, PFAM, and Prosite.

The programs described above for the assembly and analysis of full length polynucleotide  
 15 and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:14-26. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

#### IV. Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a  
 20 gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel, 1995, supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in nucleotide databases such as GenBank or LIFESEQ database (Incyte  
 25 Pharmaceuticals). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

100

30 The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1% to 2% error, and, with a product score of 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

35 The results of northern analyses are reported a percentage distribution of libraries in which

the transcript encoding CSIGP occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease or condition categories included cancer, inflammation/trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease expression are reported in Table 3.

#### V. Extension of CSIGP Encoding Polynucleotides

The full length nucleic acid sequence of SEQ ID NO:14-26 was produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing  $Mg^{2+}$ ,  $(NH_4)_2SO_4$ , and  $\beta$ -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100  $\mu$ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5  $\mu$ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5  $\mu$ l to 10  $\mu$ l aliquot of the reaction mixture

was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and  
5 sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in  
10 restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following  
15 parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulphoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer  
20 sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer).

In like manner, the nucleotide sequence of SEQ ID NO:14-26 is used to obtain 5' regulatory sequences using the procedure above, oligonucleotides designed for such extension, and an appropriate genomic library.

## 25 VI. Choice, Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:14-26 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06  
30 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 µCi of [ $\gamma$ -<sup>32</sup>P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10<sup>7</sup> counts per minute of the labeled probe is used in a typical membrane-  
35 based hybridization analysis of human genomic DNA digested with one of the following

endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature  
5 under increasingly stringent conditions up to 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. After XOMAT-AR film (Eastman Kodak, Rochester NY) is exposed to the blots to film for several hours, hybridization patterns are compared visually.

## VII. Microarrays

A chemical coupling procedure and an ink jet device can be used to synthesize array  
10 elements on the surface of a substrate. (See, e.g., Baldeschweiler, supra.) An array analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements. After hybridization, nonhybridized probes are removed and a scanner used to determine the levels  
15 and patterns of fluorescence. The degree of complementarity and the relative abundance of each probe which hybridizes to an element on the microarray may be assessed through analysis of the scanned images.

Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may comprise the elements of the microarray. Fragments suitable for hybridization can be selected  
20 using software well known in the art such as LASERGENE software (DNASTAR). Full-length cDNAs, ESTs, or fragments thereof corresponding to one of the nucleotide sequences of the present invention, or selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking followed by thermal and chemical treatments and subsequent drying. (See, e.g.,  
25 Schena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

## VIII. Complementary Polynucleotides

Sequences complementary to the CSIGP-encoding sequences, or any parts thereof, are  
30 used to detect, decrease, or inhibit expression of naturally occurring CSIGP. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of CSIGP. To inhibit transcription, a complementary oligonucleotide is designed from the most  
35 unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit

translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the CSIGP-encoding transcript.

#### IX. Expression of CSIGP

Expression and purification of CSIGP is achieved using bacterial or virus-based  
5 expression systems. For expression of CSIGP in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac* (*tac*) hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts,  
10 e.g., BL21(DE3). Antibiotic resistant bacteria express CSIGP upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of CSIGP in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding CSIGP by either homologous recombination  
15 or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E. K. et al. (1994) Proc. Natl. Acad. Sci. USA  
20 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, CSIGP is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion  
25 proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from CSIGP at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine  
30 residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch 10 and 16). Purified CSIGP obtained by these methods can be used directly in the following activity assay.

#### X. Demonstration of CSIGP Activity

CSIGP activity can be assayed in vitro by monitoring the mobilization of  $Ca^{++}$  as part of  
35 the signal transduction pathway. (See, e.g., Grynkievycz, G. et al. (1985) J. Biol. Chem.



260:3440; McColl, S. et al. (1993) J. Immunol. 150:4550-4555; and Aussel, C. et al. (1988) supra)

The assay requires preloading neutrophils or T cells with a fluorescent dye such as FURA-2 or BCECF (Universal Imaging Corp, Westchester PA) whose emission characteristics have been altered by  $\text{Ca}^{++}$  binding. When the cells are exposed to one or more activating stimuli artificially  
5 (ie, anti-CD3 antibody ligation of the T cell receptor) or physiologically (ie, by allogeneic stimulation),  $\text{Ca}^{++}$  flux takes place. This flux can be observed and quantified by assaying the cells in a fluorometer or fluorescent activated cell sorter. Measurements of  $\text{Ca}^{++}$  flux are compared between cells in their normal state and those preloaded with CSIGP.

Protein kinase activity in CSIGP is determined by measuring the phosphorylation of a  
10 protein substrate using gamma-labeled  $^{32}\text{P}$ -ATP and quantitation of the incorporated radioactivity using a radioisotope counter. CSIGP is incubated with the protein substrate,  $^{32}\text{P}$ -ATP, and an appropriate kinase buffer. The  $^{32}\text{P}$  incorporated into the product is separated from free  $^{32}\text{P}$ -ATP by electrophoresis and the incorporated  $^{32}\text{P}$  is counted. The amount of  $^{32}\text{P}$  recovered is proportional to the activity of CSIGP in the assay. A determination of the specific amino acid residue  
15 phosphorylated is made by phosphoamino acid analysis of the hydrolyzed protein.

Protein phosphatase (PP) activity in CSIGP is determined by measuring the hydrolysis of P-nitrophenyl phosphate (PNPP). CSIGP is incubated together with PNPP in HEPES buffer pH 7.5, in the presence of 0.1% b-mercaptoethanol at 37°C for 60 min. The reaction is stopped by the addition of 6 ml of 10 N NaOH and the increase in light absorbance at 410 nm resulting from the  
20 hydrolysis of PNPP is measured using a spectrophotometer. The increase in light absorbance is proportional to the activity of CSIGP in the assay.

#### **XI. Production of CSIGP Specific Antibodies**

CSIGP substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) Methods Enzymol. 182:488-495), or other purification techniques, is  
25 used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the CSIGP amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are  
30 well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)

Typically, oligopeptides 15 residues in length are synthesized using an ABI 431A Peptide Synthesizer (Perkin-Elmer) using fmoc-chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, supra.) Rabbits are immunized with the  
35 oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for

antipeptide activity by, for example, binding the peptide to plastic, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

### **XII. Purification of Naturally Occurring CSIGP Using Specific Antibodies**

Naturally occurring or recombinant CSIGP is substantially purified by immunoaffinity  
5 chromatography using antibodies specific for CSIGP. An immunoaffinity column is constructed by covalently coupling anti-CSIGP antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing CSIGP are passed over the immunoaffinity column, and the column is  
10 washed under conditions that allow the preferential absorbance of CSIGP (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/CSIGP binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and CSIGP is collected.

### **XIII. Identification of Molecules Which Interact with CSIGP**

15 CSIGP, or biologically active fragments thereof, are labeled with <sup>125</sup>I Bolton-Hunter reagent. (See, e.g., Bolton et al. (1973) Biochem. J. 133:529.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled CSIGP, washed, and any wells with labeled CSIGP complex are assayed. Data obtained using different concentrations of CSIGP are used to calculate values for the number, affinity, and association of CSIGP with the  
20 candidate molecules.

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited  
25 to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	14	016108	HUVELPB01	016108, 016624, (HUVELPB01), 970134 (MUSCNOT02), 1605858 (LUNGNOT15), 1419046 (KIDNNOT09)
2	15	640521	BRSTNOT03	640521 (BRSTNOT03)
3	16	1250171	LUNGFET03	1250171 (LUNGFET03), 260744 (HNT2RAT01), 077085 (SYNORAB01), 2790184 (COLNTUT16), SAEB01398, SAEB00499, SAEB02190, SAEB00648, SAEB00948
4	17	1911587	CONNTUT01	1911587 (CONNTUT01), 1989659 (CORPNOT02)
5	18	2079081	ISLTNOT01	2079081 (ISLTNOT01), 2631449 (COLNTUT15), 2350624 (COLSUCT01), 2568459 (HIPOAZT01), 2132860 (OVARNOT03)
6	19	2472655	THP1NOT03	2472655 (THP1NOT03), 1325950 (LPARNOT02), SAEA01014, SAEA01114, SAEA03382
7	20	2948818	KIDNFET01	2948818 (KIDNFET01), 1543592 (PROSTUT04), SAAE00176

Table 1 cont.

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
8	21	054191	FIBRNOT01	054191H1 and 054191R6 (FIBRNOT01), 483547H1, 483547R6, and 483547T6 (HNT2RAT01), 1537974R6 (SINTTUT01), 1633493H1 (COLNNOT19)
9	22	1403604	LATRTUT02	491348H1 (HNT2AGT01), 1403604H1 (LATRTUT02), 3331135T6.com (BRAIFET01), SBAA02561F1.comp, SBAA03200F1, SBAA01960F1.comp, SBAA01439F1, SBAA01304F1
10	23	1652936	PROSTUT08	467767R6 (LATRNOT01), 1551938R6 (PROSNOT06), 1652936F6 and 1652936H1 (PROSTUT08), 1817388F6 and 1817388H1 (PROSNOT20), 2822521H1 (ADRETUT06)
11	24	1710702	PROSNOT16	1474380T1 (LUNGUTUT03), 1710702H1 (PROSNOT16), 2189187H1 (PROSNOT26), 1526267F1 (UCMCL5T01), 1467104F1 (PANCTUT02)
12	25	3239149	COLAUCT01	482693H1 (HNT2RAT01), 2287788R6 (BRAINON01), 2570350T6 (HIPOAZT01), 3239149F6 and 3239149H1 (COLAUCT01), 3837574F6 (DENDTNT01), 4993747H1 (LIVRTUT11)
13	26	3315936	PROSBPT03	2501356T6 (ADRETUT05), 3315936H1 (PROSBPT03)

Table 2

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential glycosylation sites	Signature Sequence	Homologous Sequence	Analytical Methods
1	418	S359 S2 T12 S56 T91 T257 S287 S306 T402 S414 T9 S16 S43 T87 S184 S327 S334	N54 N70 N118	Y58-I293	Serine /threonine protein kinase	BLOCKS PRINTS PFAM
2	540	S100 T145 S26 T56 S100 T166 S358 S456 T462 T467 S503 S11 S30 S95 S137 S197 T280 T362 S367 S474 Y234 Y305	N460	Y165-V446	Ca2 +/calmodulin- dependent protein kinase kinase	BLOCKS PRINTS MOTIFS BLAST PFAM
3	729	T96 S348 T373 S518 S531 T682 T78 T239 T478 Y235	N42 N455 N614	W9-I238	Serine/ threonine protein kinase	BLOCKS PFAM PRINTS MOTIFS BLAST
4	313	S38 S82 S95 S97 T143 Y30	N79 N80 N172 N192	R114-S135	Protein tyrosine phosphatase	PRINTS BLAST

Table 2 cont.

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential glycosylation sites	Signature Sequence	Homologous Sequence	Analytical Methods
5	506	S114 S300 S81 S160 T162 S211 S253 S291 S335 S341 T63 S143 T144 S156 T177 S196 S363 S439 Y45 Y187	N275	SH3 domains: R441-L495	PEST phosphatase interacting protein	BLOCKS PRINTS PFAM BLAST
6	341	S39 S118 T125 S180 S110 S170 S173 S195 T299	N37 N178 N229 N263		Prolactin receptor associated protein (PRAP)	BLAST
7	898	S56 T640 S15 S107 T210 T267 S324 S366 S374 S504 T547 T592 T640 S655 T681 T756 S775 S58 S249 T437 S551 T573 S655 T726 T745 T762 S836 S858 S879	N322 N347 N389 N502 N503	F24-V277	Serine/ threonine protein kinase	BLOCKS PRINTS PFAM MOTIFS BLAST

Table 2 cont.

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential glycosylation sites	Signature Sequence	Homologous Sequence	Analytical Methods
8	336	S34 T110 S148 S311	N137 N144 N169	T175-I195 V236-T254	putative G- protein-coupled receptor	PRINTS, BLAST HMM, Motifs
9	686	T192 S312 S483 S502 S23 T584	N17 N457 N618 N642	G544-N560	GDP-GTP exchange protein	PRINTS, BLAST Motifs
10	519	S3 S77 S130 S176 S187 T196 S245 S265 T280 T290 T305 T324 S325 S351 S384 S390 T29 S33 S265 T305 S311 T453 S464 Y131 Y145	N128		GTPase-interacting protein	BLAST Motifs
11	334	S332 T186 S198 S269 T321 S90 S139 Y289	N20 N30	L267-L281	G-protein beta WD-40 repeat containing protein	PRINTS, BLAST Motifs
12	569	S91 S19 S109 S162 S376 S418 T514 S535 S536 S19 S39 T266 T288 T328 T381 T411 T451 S519	N17 N77 N416	I320-V334 M360-M374 I403-T417 V443-I457 I483-L497 I532-F546	beta-transducin repeats containing protein	PRINTS, BLAST PFAM, Motifs
13	123	S14 T107 Y44 Y70	N100	M1-N52	SAR1 family GTP-binding protein	PRINTS, BLOCKS BLAST, Motifs

Table 3

Polynucleotide SEQ ID NO:	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
14	Cardiovascular (0.194) Hematopoietic/Immune (0.194) Developmental (0.139)	Cancer (0.389) Inflammation (0.333) Cell proliferative (0.306)	pBLUESCRIPT
15	Reproductive (0.282) Nervous (0.179) Developmental (0.128)	Cancer (0.410) Cell proliferative (0.205) Inflammation (0.154)	pSPORT1
16	Reproductive (0.286) Hematopoietic/Immune (0.167) Nervous (0.119)	Cancer (0.429) Inflammation (0.310) Cell proliferative (0.214)	pINCY
17	Nervous (0.235) Reproductive (0.147) Gastrointestinal (0.118)	Cancer (0.471) Cell proliferative (0.176) Trauma (0.176)	pINCY
18	Reproductive (0.400) Gastrointestinal (0.267) Cardiovascular (0.133)	Cancer (0.533) Inflammation (0.333) Cell proliferative (0.067)	pINCY
19	Nervous (0.273) Hematopoietic/Immune (0.227) Reproductive (0.227)	Cancer (0.364) Inflammation (0.364) Cell proliferative (0.318)	pINCY
20	Hematopoietic/Immune (0.216) Reproductive (0.216) Nervous (0.157)	Cancer (0.412) Inflammation (0.294) Cell proliferative (0.216)	pINCY



Table 3 cont.

Polynucleotide SEQ ID NO:	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
21	Cardiovascular (0.217) Gastrointestinal (0.174) Nervous (0.174)	Cell proliferative (0.652) Inflammation (0.304)	pBLUESCRIPT
22	Reproductive (0.370) Nervous (0.222) Hematopoietic/Immune (0.148)	Cell proliferative (0.778) Trauma (0.148)	pINCY
23	Reproductive (0.400) Cardiovascular (0.200) Hematopoietic/Immune (0.133)	Cancer (0.533) Inflammation (0.200)	pINCY
24	Reproductive (0.241) Nervous (0.190) Cardiovascular (0.138)	Cell proliferative (0.724) Inflammation (0.138)	pINCY
25	Musculoskeletal (0.222) Nervous (0.222) Gastrointestinal (0.167)	Cell proliferative (0.555) Inflammation (0.222)	pINCY
26	Reproductive (0.750) Cardiovascular (0.250)	Cancer (0.500) Inflammation (0.500)	pINCY

0

Table 4

Polynucleotide SEQ ID NO:	Library	Library Description
14	HUVELPB01	The library was constructed using RNA isolated from HUV-EC-C (ATCC CRL 1730) cells that were stimulated with cytokine/LPS. HUV-EC-C is an endothelial cell line derived from the vein of a normal human umbilical cord. RNA was isolated from two pools of HUV-EC-C cells that had been treated with either gamma IFN and TNF-alpha or IL-1 beta and LPS.
15	BRSTNOT03	The library was constructed using RNA isolated from nontumorous breast tissue removed from a 54-year-old Caucasian female during a bilateral radical mastectomy. Pathology for the associated tumor tissue indicated residual invasive grade 3 mammary ductal adenocarcinoma. Family history included benign hypertension, hyperlipidemia, and a malignant neoplasm of the colon.
16	LUNGFET03	The library was constructed using RNA isolated from lung tissue removed from a Caucasian female fetus, who died at 20 weeks' gestation from fetal demise. Family history included bronchitis.
17	CONNTUT01	The library was constructed using RNA isolated from a soft tissue tumor removed from the clival area of the skull of a 30-year-old Caucasian female. Pathology indicated chondroid chordoma with neoplastic cells reactive for keratin. Patient history included deficiency anemia.
18	ISLTNOT01	The library was constructed using RNA isolated from pancreatic islet cells. Starting RNA was made from a pooled collection of islet cells.
19	THPINOT03	The library was constructed using RNA isolated from untreated THP-1 cells. THP-1 (ATCC TIB 202) is a human promonocyte line derived from the peripheral blood of a 1-year-old Caucasian male with acute monocytic leukemia.
20	KIDNFET01	The library was constructed using RNA isolated from kidney tissue removed from a Caucasian female fetus, who died at 17 weeks' gestation from anencephalus. Family history included gastritis.

Table 4 cont.

Polynucleotide SEQ ID NO:	Library	Library Description
21	FIBRN0T01	The library was constructed at Stratagene (STR937212), using RNA isolated from the WI38 lung fibroblast cell line, which was derived from a 3-month-old Caucasian female fetus. 2x10 <sup>6</sup> primary clones were amplified to stabilize the library for long-term storage.
22	LATRTUT02	The library was constructed using RNA isolated from a myxoma removed from the left atrium of a 43-year-old Caucasian male during annuloplasty. Pathology indicated atrial myxoma. Patient history included pulmonary insufficiency, acute myocardial infarction, atherosclerotic coronary artery disease and hyperlipidemia. Family history included benign hypertension, acute myocardial infarction, atherosclerotic coronary artery disease, and type II diabetes.
23	PROSTUT08	The library was constructed using RNA isolated from prostate tumor tissue removed from a 60-year-old Caucasian male during radical prostatectomy and regional lymph node excision. Pathology indicated an adenocarcinoma (Gleason grade 3+4). Adenofibromatous hyperplasia was also present. The patient presented with elevated prostate specific antigen (PSA). Family history included tuberculosis, cerebrovascular disease, and arteriosclerotic coronary artery disease.
24	PROSNOT16	The library was constructed using RNA isolated from diseased prostate tissue removed from a 68-year-old Caucasian male during a radical prostatectomy. Pathology indicated adenofibromatous hyperplasia. Pathology for the associated tumor tissue indicated an adenocarcinoma (Gleason grade 3+4). The patient presented with elevated prostate specific antigen (PSA). During this hospitalization, the patient was diagnosed with myasthenia gravis. Patient history included osteoarthritis, and type II diabetes. Family history included benign hypertension, acute myocardial infarction, hyperlipidemia, and arteriosclerotic coronary artery disease.

Table 4 cont.

Polynucleotide SEQ ID NO:	Library	Library Description
25	COLAUCT01	The library was constructed using RNA isolated from diseased ascending colon tissue removed from a 74-year-old Caucasian male during a multiple-segment large bowel excision with temporary ileostomy. Pathology indicated inflammatory bowel disease consistent with chronic ulcerative colitis, severe acute and chronic mucosal inflammation with erythema, ulceration, and pseudopolyp formation involving the entire colon and rectum. The sigmoid colon had an area of mild stricture formation. One diverticulum with diverticulitis was identified near this zone.
26	PROSBPT03	The library was constructed using RNA isolated from diseased prostate tissue removed from a 59-year-old Caucasian male during a radical prostatectomy and regional lymph node excision. Pathology indicated benign prostatic hyperplasia (BPH). Pathology for the associated tumor indicated adenocarcinoma, Gleason grade 3+3. The patient presented with elevated prostate specific antigen (PSA), benign hypertension, and hyperlipidemia. Family history included cerebrovascular disease, benign hypertension and prostate cancer.

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25: 3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183: 63-98; and Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLOCKS IMPROVED Searcher that matches a sequence against those in BLOCKS and PRINTS databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S and J.G. Henikoff, Nucl. Acid Res., 19:6565-72, 1991. J.G. Henikoff and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37: 417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and Probability value= 1.0E-3 or less, if applicable
PFAM	A Hidden Markov Models-based application useful for protein family search.	Krogh, A. et al. (1994) J. Mol. Biol., 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits, depending on individual protein families

Table 5 cont.

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25: 217-221.	Score= 4.0 or greater
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12: 431-439.	Score=5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch et al. <u>supra</u> ; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. A substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-13, or a fragment thereof.
- 5 2. A substantially purified variant having at least 90% amino acid identity to the amino acid sequence of claim 1.
3. An isolated and purified polynucleotide encoding the polypeptide of claim 1.
4. An isolated and purified polynucleotide variant having at least 70% polynucleotide sequence identity to the polynucleotide of claim 3.
- 10 5. An isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide of claim 3.
6. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide of claim 3.
7. A method for detecting a polynucleotide, the method comprising the steps of:
  - 15 (a) hybridizing the polynucleotide of claim 6 to at least one nucleic acid in the sample, thereby forming a hybridization complex; and
  - (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of the polynucleotide in the sample.
8. The method of claim 7 further comprising amplifying the polynucleotide prior to  
20 hybridization.
9. An isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:14-26, or a fragment thereof.
10. An isolated and purified polynucleotide variant having at least 70% polynucleotide sequence identity to the polynucleotide of claim 9.
- 25 11. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide of claim 9.
12. An expression vector comprising at least a fragment of the polynucleotide of claim 3.
13. A host cell comprising the expression vector of claim 12.
- 30 14. A method for producing a polypeptide, the method comprising the steps of:
  - a) culturing the host cell of claim 13 under conditions suitable for the expression of the polypeptide; and
  - b) recovering the polypeptide from the host cell culture.
15. A pharmaceutical composition comprising the polypeptide of claim 1 in  
35 conjunction with a suitable pharmaceutical carrier.

16. A purified antibody which specifically binds to the polypeptide of claim 1.
17. A purified agonist of the polypeptide of claim 1.
18. A purified antagonist of the polypeptide of claim 1.
19. A method for treating or preventing a disorder associated with decreased  
5 expression of CSIGP, the method comprising administering to a subject in need of such treatment  
an effective amount of the pharmaceutical composition of claim 15.
20. A method for treating or preventing a disorder associated with increased  
expression of CSIGP, the method comprising administering to a subject in need of such treatment  
an effective amount of the antagonist of claim 18.

10



## SEQUENCE LISTING

&lt;110&gt; INCYTE PHARMACEUTICALS, INC.

BANDMAN, Olga  
 HILLMAN, Jennifer L.  
 LAL, Preeti  
 YUE, Henry  
 TANG, Y. Tom  
 PATTERSON, Chandra  
 BAUGHN, Mariah R.  
 YANG, Junming

&lt;120&gt; CELL SIGNALING PROTEINS

&lt;130&gt; PF-0521 PCT

&lt;140&gt; To Be Assigned

&lt;141&gt; Herewith

&lt;150&gt; 60/085,3434; 60/098,010

&lt;151&gt; 1998-05-13; 1998-08-26

&lt;160&gt; 26

&lt;170&gt; PERL Program

&lt;210&gt; 1

&lt;211&gt; 418

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte Clone 016108

&lt;400&gt; 1

Met	Ser	Leu	Leu	Asp	Cys	Phe	Cys	Thr	Ser	Arg	Thr	Gln	Val	Glu
1				5					10					15
Ser	Leu	Arg	Pro	Glu	Lys	Gln	Ser	Glu	Thr	Ser	Ile	His	Gln	Tyr
				20					25					30
Leu	Val	Asp	Glu	Pro	Thr	Leu	Ser	Trp	Ser	Arg	Pro	Ser	Thr	Arg
				35					40					45
Ala	Ser	Glu	Val	Leu	Cys	Ser	Thr	Asn	Val	Ser	His	Tyr	Glu	Leu
				50					55					60
Gln	Val	Glu	Ile	Gly	Arg	Gly	Phe	Asp	Asn	Leu	Thr	Ser	Val	His
				65					70					75
Leu	Ala	Arg	His	Thr	Pro	Thr	Gly	Thr	Leu	Val	Thr	Ile	Lys	Ile
				80					85					90
Thr	Asn	Leu	Glu	Asn	Cys	Asn	Glu	Glu	Arg	Leu	Lys	Ala	Leu	Gln
				95					100					105
Lys	Ala	Val	Ile	Leu	Ser	His	Phe	Phe	Arg	His	Pro	Asn	Ile	Thr
				110					115					120
Thr	Tyr	Trp	Thr	Val	Phe	Thr	Val	Gly	Ser	Trp	Leu	Trp	Val	Ile
				125					130					135
Ser	Pro	Phe	Met	Ala	Tyr	Gly	Ser	Ala	Ser	Gln	Leu	Leu	Arg	Thr
				140					145					150
Tyr	Phe	Pro	Glu	Gly	Met	Ser	Glu	Thr	Leu	Ile	Arg	Asn	Ile	Leu
				155					160					165
Phe	Gly	Ala	Val	Arg	Gly	Leu	Asn	Tyr	Leu	His	Gln	Asn	Gly	Cys
				170					175					180
Ile	His	Arg	Ser	Ile	Lys	Ala	Ser	His	Ile	Leu	Ile	Ser	Gly	Asp
				185					190					195
Gly	Leu	Val	Thr	Leu	Ser	Gly	Leu	Ser	His	Leu	His	Ser	Leu	Val

	200		205		210
Lys His Gly Gln Arg His Arg Ala Val Tyr Asp Phe Pro Gln Phe					
	215		220		225
Ser Thr Ser Val Gln Pro Trp Leu Ser Pro Glu Leu Leu Arg Gln					
	230		235		240
Asp Leu His Gly Leu Tyr Val Lys Ser Asp Ile Tyr Ser Val Gly					
	245		250		255
Ile Thr Ala Cys Glu Leu Ala Ser Gly Gln Val Pro Phe Gln Asp					
	260		265		270
Met His Arg Thr Gln Met Leu Leu Gln Lys Leu Lys Gly Pro Pro					
	275		280		285
Tyr Ser Pro Leu Asp Ile Ser Ile Phe Pro Gln Ser Glu Ser Arg					
	290		295		300
Met Lys Asn Ser Gln Ser Gly Val Asp Ser Gly Ile Gly Glu Ser					
	305		310		315
Val Leu Val Ser Ser Gly Thr His Thr Val Asn Ser Asp Arg Leu					
	320		325		330
His Thr Pro Ser Ser Lys Thr Phe Ser Pro Ala Phe Phe Ser Leu					
	335		340		345
Val Gln Leu Cys Leu Gln Gln Asp Pro Glu Lys Arg Pro Ser Ala					
	350		355		360
Ser Ser Leu Leu Ser His Val Phe Phe Lys Gln Met Lys Glu Glu					
	365		370		375
Ser Gln Asp Ser Ile Leu Ser Leu Leu Pro Pro Ala Tyr Asn Lys					
	380		385		390
Pro Ser Ile Ser Leu Pro Pro Val Leu Pro Trp Thr Glu Pro Glu					
	395		400		405
Cys Asp Phe Pro Asp Glu Lys Asp Ser Tyr Trp Glu Phe					
	410		415		

&lt;210&gt; 2

&lt;211&gt; 540

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte Clone 640521

&lt;400&gt; 2

Met Ser Ser Cys Val Ser Ser Gln Pro Ser Ser Asn Arg Ala Ala		
1 5 10 15		
Pro Gln Asp Glu Leu Gly Gly Arg Gly Ser Ser Ser Ser Glu Ser		
20 25 30		
Gln Lys Pro Cys Glu Ala Leu Arg Gly Leu Ser Ser Leu Ser Ile		
35 40 45		
His Leu Gly Met Glu Ser Phe Ile Val Val Thr Glu Cys Glu Pro		
50 55 60		
Gly Cys Ala Val Asp Leu Gly Leu Ala Arg Asp Arg Pro Leu Glu		
65 70 75		
Ala Asp Gly Gln Glu Val Pro Leu Asp Ser Ser Gly Ser Gln Ala		
80 85 90		
Arg Pro His Leu Ser Gly Arg Lys Leu Ser Leu Gln Glu Arg Ser		
95 100 105		
Gln Gly Gly Leu Ala Ala Gly Gly Ser Leu Asp Met Asn Gly Arg		
110 115 120		
Cys Ile Cys Pro Ser Leu Pro Tyr Ser Pro Val Ser Ser Pro Gln		
125 130 135		
Ser Ser Pro Arg Leu Pro Arg Arg Pro Thr Val Glu Ser His His		
140 145 150		
Val Ser Ile Thr Gly Met Gln Asp Cys Val Gln Leu Asn Gln Tyr		

Thr Leu Lys Asp	155	Ile Gly Lys Gly	160	Ser Tyr Gly Val Val	165
	170		175		180
Leu Ala Tyr Asn	185	Glu Asn Asp Asn Thr	190	Tyr Tyr Ala Met Lys	195
	200		205		210
Leu Ser Lys Lys	215	Lys Leu Ile Arg Gln	220	Ala Gly Phe Pro Arg	225
	230		235		240
Pro Pro Pro Arg	245	Gly Thr Arg Pro Ala	250	Pro Gly Gly Cys Ile	255
	260		265		270
Pro Arg Gly Pro	275	Ile Glu Gln Val Tyr	280	Gln Glu Ile Ala Ile	285
	290		295		300
Lys Lys Leu Asp	305	His Pro Asn Val Val	310	Lys Leu Val Glu Val	315
	320		325		330
Asp Asp Pro Asn	335	Glu Asp His Leu Tyr	340	Met Val Phe Glu Leu	345
	350		355		360
Asn Gln Gly Pro	365	Val Met Glu Val Pro	370	Thr Leu Lys Pro Leu	375
	380		385		390
Glu Asp Gln Ala	395	Arg Phe Tyr Phe Gln	400	Asp Leu Ile Lys Gly	405
	410		415		420
Glu Tyr Leu His	425	Tyr Gln Lys Ile Ile	430	His Arg Asp Ile Lys	435
	440		445		450
Ser Asn Leu Leu	455	Val Gly Glu Asp Gly	460	His Ile Lys Ile Ala	465
	470		475		480
Phe Gly Val Ser	485	Asn Glu Phe Lys Gly	490	Ser Asp Ala Leu Leu	495
	500		505		510
Asn Thr Val Gly	515	Thr Pro Ala Phe Met	520	Ala Pro Glu Ser Leu	525
	530		535		540
Glu Thr Arg Lys		Ile Phe Ser Gly Lys		Ala Leu Asp Val Trp	
Met Gly Val Thr		Leu Tyr Cys Phe Val		Phe Gly Gln Cys Pro	
Met Asp Glu Arg		Ile Met Cys Leu His		Ser Lys Ile Lys Ser	
Ala Leu Glu Phe		Pro Asp Gln Pro Asp		Ile Ala Glu Asp Leu	
Asp Leu Ile Thr		Arg Met Leu Asp Lys		Asn Pro Glu Ser Arg	
Val Val Pro Glu		Ile Lys Leu His Pro		Trp Val Thr Arg His	
Ala Glu Pro Leu		Pro Ser Glu Asp Glu		Asn Cys Thr Leu Val	
Val Thr Glu Glu		Glu Val Glu Asn Ser		Val Lys His Ile Pro	
Leu Ala Thr Val		Ile Leu Val Lys Thr		Met Ile Arg Lys Arg	
Phe Gly Asn Pro		Phe Glu Gly Ser Arg		Arg Glu Glu Arg Ser	
Ser Ala Pro Gly		Asn Leu Leu Thr Lys		Gln Gly Ser Glu Asp	
Leu Gln Gly Thr		Asp Pro Pro Pro Val		Gly Glu Glu Glu Val	

&lt;210&gt; 3

&lt;211&gt; 729

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte Clone 1250171

&lt;400&gt; 3

```

Met Gln Ser Thr Ser Asn His Leu Trp Leu Leu Ser Asp Ile Leu
 1          5          10          15
Gly Gln Gly Ala Thr Ala Asn Val Phe Arg Gly Arg His Lys Lys
 20          25          30
Thr Gly Asp Leu Phe Ala Ile Lys Val Phe Asn Asn Ile Ser Phe
 35          40          45
Leu Arg Pro Val Asp Val Gln Met Arg Glu Phe Glu Val Leu Lys
 50          55          60
Lys Leu Asn His Lys Asn Ile Val Lys Leu Phe Ala Ile Glu Glu
 65          70          75
Glu Thr Thr Thr Arg His Lys Val Leu Ile Met Glu Phe Cys Pro
 80          85          90
Cys Gly Ser Leu Tyr Thr Val Leu Glu Glu Pro Ser Asn Ala Tyr
 95          100          105
Gly Leu Pro Glu Ser Glu Phe Leu Ile Val Leu Arg Asp Val Val
 110          115          120
Gly Gly Met Asn His Leu Arg Glu Asn Gly Ile Val His Arg Asp
 125          130          135
Ile Lys Pro Gly Asn Ile Met Arg Val Ile Gly Glu Asp Gly Gln
 140          145          150
Ser Val Tyr Lys Leu Thr Asp Phe Gly Ala Ala Arg Glu Leu Glu
 155          160          165
Asp Asp Glu Gln Phe Val Ser Leu Tyr Gly Thr Glu Glu Tyr Leu
 170          175          180
His Pro Asp Met Tyr Glu Arg Ala Val Leu Arg Lys Asp His Gln
 185          190          195
Lys Lys Tyr Gly Ala Thr Val Asp Leu Trp Ser Ile Gly Val Thr
 200          205          210
Phe Tyr His Ala Ala Thr Gly Ser Leu Pro Phe Arg Pro Phe Glu
 215          220          225
Gly Pro Arg Arg Asn Lys Glu Val Met Tyr Lys Ile Ile Thr Gly
 230          235          240
Lys Pro Ser Gly Ala Ile Ser Gly Val Gln Lys Ala Glu Asn Gly
 245          250          255
Pro Ile Asp Trp Ser Gly Asp Met Pro Val Ser Cys Ser Leu Ser
 260          265          270
Arg Gly Leu Gln Val Leu Leu Thr Pro Val Leu Ala Asn Ile Leu
 275          280          285
Glu Ala Asp Gln Glu Lys Cys Trp Gly Phe Asp Gln Phe Phe Ala
 290          295          300
Glu Thr Ser Asp Ile Leu His Arg Met Val Ile His Val Phe Ser
 305          310          315
Leu Gln Gln Met Thr Ala His Lys Ile Tyr Ile His Ser Tyr Asn
 320          325          330
Thr Ala Thr Ile Phe His Glu Leu Val Tyr Lys Gln Thr Lys Ile
 335          340          345
Ile Ser Ser Asn Gln Glu Leu Ile Tyr Glu Gly Arg Arg Leu Val
 350          355          360
Leu Glu Pro Gly Arg Leu Ala Gln His Phe Pro Lys Thr Thr Glu
 365          370          375
Glu Asn Pro Ile Phe Val Val Ser Arg Glu Pro Leu Asn Thr Ile
 380          385          390
Gly Leu Ile Tyr Glu Lys Ile Ser Leu Pro Lys Val His Pro Arg
 395          400          405
Tyr Asp Leu Asp Gly Asp Ala Ser Met Ala Lys Ala Ile Thr Gly
 410          415          420
Val Val Cys Tyr Ala Cys Arg Ile Ala Ser Thr Leu Leu Leu Tyr
 425          430          435
Gln Glu Leu Met Arg Lys Gly Ile Arg Trp Leu Ile Glu Leu Ile
 440          445          450
Lys Asp Asp Tyr Asn Glu Thr Val His Lys Lys Thr Glu Val Val
 455          460          465
Ile Thr Leu Asp Phe Cys Ile Arg Asn Ile Glu Lys Thr Val Lys

```

	470		475		480
Val Tyr Glu Lys	Leu Met Lys Ile Asn	Leu Glu Ala Ala Glu	Leu		
	485		490		495
Gly Glu Ile Ser	Asp Ile His Thr Lys	Leu Leu Arg Leu Ser	Ser		
	500		505		510
Ser Gln Gly Thr	Ile Glu Thr Ser Leu	Gln Asp Ile Asp Ser	Arg		
	515		520		525
Leu Ser Pro Gly	Gly Ser Leu Ala Asp	Ala Trp Ala His Gln	Glu		
	530		535		540
Gly Thr His Pro	Lys Asp Arg Asn Val	Glu Lys Leu Gln Val	Leu		
	545		550		555
Leu Asn Cys Met	Thr Glu Ile Tyr Tyr	Gln Phe Lys Lys Asp	Lys		
	560		565		570
Ala Glu Arg Arg	Glu Ala Tyr Asn Glu	Gln Ile His Lys	Phe		
	575		580		585
Asp Lys Gln Lys	Leu Tyr Tyr His Ala	Thr Lys Ala Met Thr	His		
	590		595		600
Phe Thr Asp Glu	Cys Val Lys Lys Tyr	Glu Ala Phe Leu Asn	Lys		
	605		610		615
Ser Glu Glu Trp	Ile Arg Lys Met Leu	His Leu Arg Lys Gln	Leu		
	620		625		630
Leu Ser Leu Thr	Asn Gln Cys Phe Asp	Ile Glu Glu Glu Val	Ser		
	635		640		645
Lys Tyr Gln Glu	Tyr Thr Asn Glu Leu	Gln Glu Thr Leu Pro	Gln		
	650		655		660
Lys Met Phe Thr	Ala Ser Ser Gly Ile	Lys His Thr Met Thr	Pro		
	665		670		675
Ile Tyr Pro Ser	Ser Asn Thr Leu Val	Glu Met Thr Leu Gly	Met		
	680		685		690
Lys Lys Leu Lys	Glu Glu Met Glu Gly	Val Val Lys Glu Leu	Ala		
	695		700		705
Glu Asn Asn His	Ile Leu Glu Arg Phe	Gly Ser Leu Thr Met	Asp		
	710		715		720
Gly Gly Leu Arg	Asn Val Asp Cys Leu				
	725				

<210> 4  
 <211> 313  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte Clone 1911587

<400> 4  
 Met Pro Gly Leu Leu Leu Cys Glu Pro Thr Glu Leu Tyr Asn Ile  
 1 5 10 15  
 Leu Asn Gln Ala Thr Lys Leu Ser Arg Leu Thr Asp Pro Asn Tyr  
 20 25 30  
 Leu Cys Leu Leu Asp Val Arg Ser Lys Trp Glu Tyr Asp Glu Ser  
 35 40 45  
 His Val Ile Thr Ala Leu Arg Val Lys Lys Lys Asn Asn Glu Tyr  
 50 55 60  
 Leu Leu Pro Glu Ser Val Asp Leu Glu Cys Val Lys Tyr Cys Val  
 65 70 75  
 Val Tyr Asp Asn Asn Ser Ser Thr Leu Glu Ile Leu Leu Lys Asp  
 80 85 90  
 Asp Asp Asp Asp Ser Asp Ser Asp Gly Asp Gly Lys Asp Leu Val  
 95 100 105  
 Pro Gln Ala Ala Ile Glu Tyr Gly Arg Ile Leu Thr Arg Leu Thr

				110					115				120	
His	His	Pro	Val	Tyr	Ile	Leu	Lys	Gly	Gly	Tyr	Glu	Arg	Phe	Ser
				125					130					135
Gly	Thr	Tyr	His	Phe	Leu	Arg	Thr	Gln	Lys	Ile	Ile	Trp	Met	Pro
				140					145					150
Gln	Glu	Leu	Asp	Ala	Phe	Gln	Pro	Tyr	Pro	Ile	Glu	Ile	Val	Pro
				155					160					165
Gly	Lys	Val	Phe	Val	Gly	Asn	Phe	Ser	Gln	Ala	Cys	Asp	Pro	Lys
				170					175					180
Ile	Gln	Lys	Asp	Leu	Lys	Ile	Lys	Ala	His	Val	Asn	Val	Ser	Met
				185					190					195
Asp	Thr	Gly	Pro	Phe	Phe	Ala	Gly	Asp	Ala	Asp	Arg	Leu	Leu	His
				200					205					210
Ile	Arg	Ile	Glu	Asp	Ser	Pro	Glu	Ala	Gln	Ile	Leu	Pro	Phe	Leu
				215					220					225
Arg	His	Met	Cys	His	Phe	Ile	Glu	Ile	His	His	His	Leu	Gly	Ser
				230					235					240
Val	Ile	Leu	Ile	Phe	Ser	Thr	Gln	Gly	Ile	Ser	Arg	Ser	Cys	Ala
				245					250					255
Ala	Ile	Ile	Ala	Tyr	Leu	Met	His	Ser	Asn	Glu	Gln	Thr	Leu	Gln
				260					265					270
Arg	Ser	Trp	Ala	Tyr	Val	Lys	Lys	Cys	Lys	Asn	Asn	Met	Cys	Pro
				275					280					285
Asn	Arg	Gly	Leu	Val	Ser	Gln	Leu	Leu	Glu	Trp	Glu	Lys	Thr	Ile
				290					295					300
Leu	Gly	Asp	Ser	Ile	Thr	Asn	Ile	Met	Asp	Pro	Leu	Tyr		
				305					310					

&lt;210&gt; 5

&lt;211&gt; 506

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte Clone 2079081

&lt;400&gt; 5

Met	Arg	Asp	Pro	Leu	Thr	Asp	Cys	Pro	Tyr	Asn	Lys	Val	Tyr	Lys
1				5					10					15
Asn	Leu	Lys	Glu	Phe	Ser	Gln	Asn	Gly	Glu	Asn	Phe	Cys	Lys	Gln
				20					25					30
Val	Thr	Ser	Val	Leu	Gln	Gln	Arg	Ala	Asn	Leu	Glu	Ile	Ser	Tyr
				35					40					45
Ala	Lys	Gly	Leu	Gln	Lys	Leu	Ala	Ser	Lys	Leu	Ser	Lys	Ala	Leu
				50					55					60
Gln	Asn	Thr	Arg	Lys	Ser	Cys	Val	Ser	Ser	Ala	Trp	Ala	Trp	Ala
				65					70					75
Ser	Glu	Gly	Met	Lys	Ser	Thr	Ala	Asp	Leu	His	Gln	Lys	Leu	Gly
				80					85					90
Lys	Ala	Ile	Glu	Leu	Glu	Ala	Ile	Lys	Pro	Thr	Tyr	Gln	Val	Leu
				95					100					105
Asn	Val	Gln	Glu	Lys	Lys	Arg	Lys	Ser	Leu	Asp	Asn	Glu	Val	Glu
				110					115					120
Lys	Thr	Ala	Asn	Leu	Val	Ile	Ser	Asn	Trp	Asn	Gln	Gln	Ile	Lys
				125					130					135
Ala	Lys	Lys	Lys	Leu	Met	Val	Ser	Thr	Lys	Lys	His	Glu	Ala	Leu
				140					145					150
Phe	Gln	Leu	Val	Glu	Ser	Ser	Lys	Gln	Ser	Met	Thr	Glu	Lys	Glu
				155					160					165
Lys	Arg	Lys	Leu	Leu	Asn	Lys	Leu	Thr	Lys	Ser	Thr	Glu	Lys	Leu

	170		175		180
Glu Lys Glu Asp	Glu Asn Tyr Tyr Gln	Lys Asn Met Ala Gly Tyr			
	185		190		195
Ser Thr Arg Leu	Lys Trp Glu Asn Thr	Leu Glu Asn Cys Tyr Gln			
	200		205		210
Ser Ile Leu Glu	Leu Glu Lys Glu Arg	Ile Gln Leu Leu Cys Asn			
	215		220		225
Asn Leu Asn Gln	Tyr Ser Gln His Ile	Ser Leu Phe Gly Gln Thr			
	230		235		240
Leu Thr Thr Cys	His Thr Gln Ile His	Cys Ala Ile Ser Lys Ile			
	245		250		255
Asp Ile Glu Lys	Asp Ile Gln Ala Val	Met Glu Glu Thr Ala Ile			
	260		265		270
Leu Ser Thr Glu	Asn Lys Ser Glu Phe	Leu Leu Thr Asp Tyr Phe			
	275		280		285
Glu Glu Asp Pro	Asn Ser Ala Met Asp	Lys Glu Arg Arg Lys Ser			
	290		295		300
Leu Leu Lys Pro	Lys Leu Leu Arg Leu	Gln Arg Asp Ile Glu Lys			
	305		310		315
Ala Ser Lys Asp	Lys Glu Gly Leu Glu	Arg Met Leu Lys Thr Tyr			
	320		325		330
Ser Ser Thr Ser	Ser Phe Ser Asp Ala	Lys Ser Gln Lys Asp Thr			
	335		340		345
Ala Ala Leu Met	Asp Glu Asn Asn Leu	Lys Leu Asp Leu Leu Glu			
	350		355		360
Ala Asn Ser Tyr	Lys Leu Ser Ser Met	Leu Ala Glu Leu Glu Gln			
	365		370		375
Arg Pro Gln Pro	Ser His Pro Cys Ser	Asn Ser Ile Phe Arg Trp			
	380		385		390
Arg Glu Lys Glu	His Thr His Ser Tyr	Val Lys Ile Ser Arg Pro			
	395		400		405
Phe Leu Met Lys	Arg Leu Glu Asn Ile	Val Ser Lys Ala Ser Ser			
	410		415		420
Gly Gly Gln Ser	Asn Pro Gly Ser Ser	Thr Pro Ala Pro Gly Ala			
	425		430		435
Ala Gln Leu Ser	Ser Arg Leu Cys Lys	Ala Leu Tyr Ser Phe Gln			
	440		445		450
Ala Arg Gln Asp	Asp Glu Leu Asn Leu	Glu Lys Gly Asp Ile Val			
	455		460		465
Ile Ile His Glu	Lys Lys Glu Glu Gly	Trp Trp Phe Gly Ser Leu			
	470		475		480
Asn Gly Lys Lys	Gly His Phe Pro Ala	Ala Tyr Val Glu Glu Leu			
	485		490		495
Pro Ser Asn Ala	Gly Asn Thr Ala Thr	Lys Ala			
	500		505		

&lt;210&gt; 6

&lt;211&gt; 341

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte Clone 2472655

&lt;400&gt; 6

Met Arg Lys Val	Val Leu Ile Thr Gly	Ala Ser Ser Gly Ile Gly
1	5	10
Leu Ala Leu Cys	Lys Arg Leu Leu Ala	Glu Asp Asp Glu Leu His
20	25	30
Leu Cys Leu Ala	Cys Arg Asn Met Ser	Lys Ala Glu Ala Val Cys

	35		40		45									
Ala	Ala	Leu	Leu	Ala	Ser	His	Pro	Thr	Ala	Glu	Val	Thr	Ile	Val
	50									55				60
Gln	Val	Asp	Val	Ser	Asn	Leu	Gln	Ser	Val	Phe	Arg	Ala	Ser	Lys
	65									70				75
Glu	Leu	Lys	Gln	Arg	Phe	Gln	Arg	Leu	Asp	Cys	Ile	Tyr	Leu	Asn
	80									85				90
Ala	Gly	Ile	Met	Pro	Asn	Pro	Gln	Leu	Asn	Ile	Lys	Ala	Leu	Phe
	95									100				105
Phe	Gly	Leu	Phe	Ser	Arg	Lys	Val	Ile	His	Met	Phe	Ser	Thr	Ala
	110									115				120
Glu	Gly	Leu	Leu	Thr	Gln	Gly	Asp	Lys	Ile	Thr	Ala	Asp	Gly	Leu
	125									130				135
Gln	Glu	Val	Phe	Glu	Thr	Asn	Val	Phe	Gly	His	Phe	Ile	Leu	Ile
	140									145				150
Arg	Glu	Leu	Glu	Pro	Leu	Leu	Cys	His	Ser	Asp	Asn	Pro	Ser	Gln
	155									160				165
Leu	Ile	Trp	Thr	Ser	Ser	Arg	Ser	Ala	Arg	Lys	Ser	Asn	Phe	Ser
	170									175				180
Leu	Glu	Asp	Phe	Gln	His	Ser	Lys	Gly	Lys	Glu	Pro	Tyr	Ser	Ser
	185									190				195
Ser	Lys	Tyr	Ala	Thr	Asp	Leu	Leu	Ser	Val	Ala	Leu	Asn	Arg	Asn
	200									205				210
Phe	Asn	Gln	Gln	Gly	Leu	Tyr	Ser	Asn	Val	Ala	Cys	Pro	Gly	Thr
	215									220				225
Ala	Leu	Thr	Asn	Leu	Thr	Tyr	Gly	Ile	Leu	Pro	Pro	Phe	Ile	Trp
	230									235				240
Thr	Leu	Leu	Met	Pro	Ala	Ile	Leu	Leu	Leu	Arg	Phe	Phe	Ala	Asn
	245									250				255
Ala	Phe	Thr	Leu	Thr	Pro	Tyr	Asn	Gly	Thr	Glu	Ala	Leu	Val	Trp
	260									265				270
Leu	Phe	His	Gln	Lys	Pro	Glu	Ser	Leu	Asn	Pro	Leu	Ile	Lys	Tyr
	275									280				285
Leu	Ser	Ala	Thr	Thr	Gly	Phe	Gly	Arg	Asn	Tyr	Ile	Met	Thr	Gln
	290									295				300
Lys	Met	Asp	Leu	Asp	Glu	Asp	Thr	Ala	Glu	Lys	Phe	Tyr	Gln	Lys
	305									310				315
Leu	Leu	Glu	Leu	Glu	Lys	His	Ile	Arg	Val	Thr	Ile	Gln	Lys	Thr
	320									325				330
Asp	Asn	Gln	Ala	Arg	Leu	Ser	Gly	Ser	Cys	Leu				
	335									340				

&lt;210&gt; 7

&lt;211&gt; 898

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte Clone 2948818

&lt;400&gt; 7

Met	Arg	Lys	Gly	Val	Leu	Lys	Asp	Pro	Glu	Ile	Ala	Asp	Leu	Ser
1				5					10					15
Tyr	Lys	Asp	Asp	Pro	Glu	Glu	Leu	Phe	Ile	Gly	Leu	His	Glu	Ile
				20					25					30
Gly	His	Gly	Ser	Phe	Gly	Ala	Val	Tyr	Phe	Ala	Thr	Asn	Ala	His
				35					40					45
Thr	Ser	Glu	Val	Val	Ala	Ile	Lys	Lys	Met	Ser	Tyr	Ser	Gly	Lys
				50					55					60
Gln	Thr	His	Glu	Lys	Trp	Gln	Asp	Ile	Leu	Lys	Glu	Val	Lys	Phe



				65					70					75
Leu	Arg	Gln	Leu	Lys	His	Pro	Asn	Thr	Ile	Glu	Tyr	Lys	Gly	Cys
				80					85					90
Tyr	Leu	Lys	Glu	His	Thr	Ala	Trp	Leu	Val	Met	Glu	Tyr	Cys	Leu
				95					100					105
Gly	Ser	Ala	Ser	Asp	Leu	Leu	Glu	Val	His	Lys	Lys	Pro	Leu	Gln
				110					115					120
Glu	Val	Glu	Ile	Ala	Ala	Ile	Thr	His	Gly	Ala	Leu	His	Gly	Leu
				125					130					135
Ala	Tyr	Leu	His	Ser	His	Ala	Leu	Ile	His	Arg	Asp	Ile	Lys	Ala
				140					145					150
Gly	Asn	Ile	Leu	Leu	Thr	Glu	Pro	Gly	Gln	Val	Lys	Leu	Ala	Asp
				155					160					165
Phe	Gly	Ser	Ala	Ser	Met	Ala	Ser	Pro	Ala	Asn	Ser	Phe	Val	Gly
				170					175					180
Thr	Pro	Tyr	Trp	Met	Ala	Pro	Glu	Val	Ile	Leu	Ala	Met	Asp	Glu
				185					190					195
Gly	Gln	Tyr	Asp	Gly	Lys	Val	Asp	Ile	Trp	Ser	Leu	Gly	Ile	Thr
				200					205					210
Cys	Ile	Glu	Leu	Ala	Glu	Arg	Lys	Pro	Pro	Leu	Phe	Asn	Met	Asn
				215					220					225
Ala	Met	Ser	Ala	Leu	Tyr	His	Ile	Ala	Gln	Asn	Asp	Ser	Pro	Thr
				230					235					240
Leu	Gln	Ser	Asn	Glu	Trp	Thr	Asp	Ser	Phe	Arg	Arg	Phe	Val	Asp
				245					250					255
Tyr	Cys	Leu	Gln	Lys	Ile	Pro	Gln	Glu	Arg	Pro	Thr	Ser	Ala	Glu
				260					265					270
Leu	Leu	Arg	His	Asp	Phe	Val	Arg	Arg	Asp	Arg	Pro	Leu	Arg	Val
				275					280					285
Leu	Ile	Asp	Leu	Ile	Gln	Arg	Thr	Lys	Asp	Ala	Val	Arg	Glu	Leu
				290					295					300
Asp	Asn	Leu	Gln	Tyr	Arg	Lys	Met	Lys	Lys	Ile	Leu	Phe	Gln	Glu
				305					310					315
Thr	Arg	Asn	Gly	Pro	Leu	Asn	Glu	Ser	Gln	Glu	Asp	Glu	Glu	Asp
				320					325					330
Ser	Glu	His	Gly	Thr	Ser	Leu	Asn	Arg	Glu	Met	Asp	Ser	Leu	Gly
				335					340					345
Ser	Asn	His	Ser	Ile	Pro	Ser	Met	Ser	Val	Ser	Thr	Gly	Ser	Gln
				350					355					360
Ser	Ser	Ser	Val	Asn	Ser	Met	Gln	Glu	Val	Met	Asp	Glu	Ser	Ser
				365					370					375
Ser	Glu	Leu	Val	Met	Met	His	Asp	Asp	Glu	Ser	Thr	Ile	Asn	Ser
				380					385					390
Ser	Ser	Ser	Val	Val	His	Lys	Lys	Asp	His	Val	Phe	Ile	Arg	Asp
				395					400					405
Glu	Ala	Gly	His	Gly	Asp	Pro	Arg	Pro	Glu	Pro	Arg	Pro	Thr	Gln
				410					415					420
Ser	Val	Gln	Ser	Gln	Ala	Leu	His	Tyr	Arg	Asn	Arg	Glu	Arg	Phe
				425					430					435
Ala	Thr	Ile	Lys	Ser	Ala	Ser	Leu	Val	Thr	Arg	Gln	Ile	His	Glu
				440					445					450
His	Glu	Gln	Glu	Asn	Glu	Leu	Arg	Glu	Gln	Met	Ser	Gly	Tyr	Lys
				455					460					465
Arg	Met	Arg	Arg	Gln	His	Gln	Lys	Gln	Leu	Ile	Ala	Leu	Glu	Asn
				470					475					480
Lys	Leu	Lys	Ala	Glu	Met	Asp	Glu	His	Arg	Leu	Lys	Leu	Gln	Lys
				485					490					495
Glu	Val	Glu	Thr	His	Ala	Asn	Asn	Ser	Ser	Ile	Glu	Leu	Glu	Lys
				500					505					510
Leu	Ala	Lys	Lys	Gln	Val	Ala	Ile	Ile	Glu	Lys	Glu	Ala	Lys	Val
				515					520					525
Ala	Ala	Ala	Asp	Glu	Lys	Lys	Phe	Gln	Gln	Gln	Ile	Leu	Ala	Gln
				530					535					540
Gln	Lys	Lys	Asp	Leu	Thr	Thr	Phe	Leu	Glu	Ser	Gln	Lys	Lys	Gln

	545		550		555
Tyr Lys Ile Cys	Lys Glu Lys Ile Lys	Glu Glu Met Asn Glu	Asp		
	560		565		570
His Ser Thr Pro	Lys Lys Glu Lys Gln	Glu Arg Ile Ser Lys	His		
	575		580		585
Lys Glu Asn Leu	Gln His Thr Gln Ala	Glu Glu Glu Ala His	Leu		
	590		595		600
Leu Thr Gln Gln	Arg Leu Tyr Tyr Asp	Lys Asn Cys Arg Phe	Phe		
	605		610		615
Lys Arg Lys Ile	Met Ile Lys Arg His	Glu Val Glu Gln Gln	Asn		
	620		625		630
Ile Arg Glu Glu	Leu Asn Lys Lys Arg	Thr Gln Lys Glu Met	Glu		
	635		640		645
His Ala Met Leu	Ile Arg His Asp Glu	Ser Thr Arg Glu Leu	Glu		
	650		655		660
Tyr Arg Gln Leu	His Thr Leu Gln Lys	Leu Arg Met Asp Leu	Ile		
	665		670		675
Arg Leu Gln His	Gln Thr Glu Leu Glu	Asn Gln Leu Glu Tyr	Asn		
	680		685		690
Lys Arg Arg Glu	Arg Glu Leu His Arg	Lys His Val Met Glu	Leu		
	695		700		705
Arg Gln Gln Pro	Lys Asn Leu Lys Ala	Met Glu Met Gln Ile	Lys		
	710		715		720
Lys Gln Phe Gln	Asp Thr Cys Lys Val	Gln Thr Lys Gln Tyr	Lys		
	725		730		735
Ala Leu Lys Asn	His Gln Leu Glu Val	Thr Pro Lys Asn Glu	His		
	740		745		750
Lys Thr Ile Leu	Lys Thr Leu Lys Asp	Glu Gln Thr Arg Lys	Leu		
	755		760		765
Ala Ile Leu Ala	Glu Gln Tyr Glu Gln	Ser Ile Asn Glu Met	Met		
	770		775		780
Ala Ser Gln Ala	Leu Arg Leu Asp Glu	Ala Gln Glu Ala Glu	Cys		
	785		790		795
Gln Ala Leu Arg	Leu Gln Leu Gln Gln	Glu Met Glu Leu Leu	Asn		
	800		805		810
Ala Tyr Gln Ser	Lys Ile Lys Met Gln	Thr Glu Ala Gln His	Glu		
	815		820		825
Arg Glu Leu Gln	Lys Leu Glu Gln Arg	Val Ser Leu Arg Arg	Ala		
	830		835		840
His Leu Glu Gln	Lys Ile Glu Glu Glu	Leu Ala Ala Leu Gln	Lys		
	845		850		855
Glu Arg Ser Glu	Arg Ile Lys Asn Leu	Leu Glu Arg Gln Glu	Arg		
	860		865		870
Glu Ile Glu Thr	Phe Asp Met Glu Ser	Leu Arg Met Gly Phe	Gly		
	875		880		885
Asn Leu Val Thr	Leu Asp Phe Pro Lys	Glu Asp Tyr Arg			
	890		895		

&lt;210&gt; 8

&lt;211&gt; 336

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte Clone 054191

&lt;400&gt; 8

Met	Ala	Thr	Leu	Ser	Val	Ile	Gly	Ser	Ser	Ser	Leu	Ile	Ala	Tyr
1				5				10						15
Ala	Val	Phe	His	Asn	Ile	Gln	Lys	Ser	Pro	Glu	Ile	Arg	Pro	Leu

	20		25		30
Phe Tyr Leu Ser	Phe Cys Asp Leu Leu Leu	Gly Leu Cys Trp	Leu		
	35		40		45
Thr Glu Thr Leu	Leu Tyr Gly Ala Ser Val	Ala Asn Lys Asp	Ile		
	50		55		60
Ile Cys Tyr Asn	Leu Gln Ala Val Gly Gln	Ile Phe Tyr Ile	Ser		
	65		70		75
Ser Phe Leu Tyr	Thr Val Asn Tyr Ile Trp	Tyr Leu Tyr Thr	Glu		
	80		85		90
Leu Arg Met Lys	His Thr Gln Ser Gly Gln	Ser Thr Ser Pro	Leu		
	95		100		105
Val Ile Asp Tyr	Thr Cys Arg Val Gly Gln	Met Ala Phe Val	Phe		
	110		115		120
Ser Ser Leu Ile	Pro Leu Leu Leu Met Thr	Pro Val Phe Cys	Leu		
	125		130		135
Gly Asn Thr Ser	Glu Cys Phe Gln Asn Phe	Ser Gln Ser His	Lys		
	140		145		150
Cys Ile Leu Met	His Ser Pro Pro Ser Ala	Met Ala Glu Leu	Pro		
	155		160		165
Pro Ser Ala Asn	Thr Ser Val Cys Ser Thr	Leu Tyr Phe Tyr	Gly		
	170		175		180
Ile Ala Ile Phe	Leu Gly Ser Phe Val Leu	Ser Leu Leu Thr	Ile		
	185		190		195
Met Val Leu Leu	Ile Arg Ala Gln Thr Leu	Tyr Lys Lys Phe	Val		
	200		205		210
Lys Ser Thr Gly	Phe Leu Gly Ser Glu Gln	Trp Ala Val Ile	His		
	215		220		225
Ile Val Asp Gln	Arg Val Arg Phe Tyr Pro	Val Ala Phe Phe	Cys		
	230		235		240
Cys Trp Gly Pro	Ala Val Ile Leu Met Ile	Ile Lys Leu Thr	Lys		
	245		250		255
Pro Gln Asp Thr	Lys Leu His Met Ala Leu	Tyr Val Leu Gln	Ala		
	260		265		270
Leu Thr Ala Thr	Ser Gln Gly Leu Leu Asn	Cys Gly Val Tyr	Gly		
	275		280		285
Trp Thr Gln His	Lys Phe His Gln Leu Lys	Gln Glu Ala Arg	Arg		
	290		295		300
Asp Ala Asp Thr	Gln Thr Pro Leu Leu Cys	Ser Gln Lys Arg	Phe		
	305		310		315
Tyr Ser Arg Gly	Leu Asn Ser Leu Glu Ser	Thr Leu Thr Phe	Pro		
	320		325		330
Ala Ser Thr Ser	Thr Ile				
	335				

&lt;210&gt; 9

&lt;211&gt; 686

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte Clone 1403604

&lt;400&gt; 9

Met Gly Pro Arg	Ser Arg Glu Arg Arg Ala	Gly Ala Val Gln	Asn
1	5	10	15
Thr Asn Asp Ser	Ser Ala Leu Ser Lys Arg	Ser Leu Ala Ala	Arg
	20	25	30
Gly Tyr Val Gln	Asp Pro Phe Ala Ala Leu	Leu Val Pro Gly	Ala
	35	40	45
Ala Arg Arg Ala	Pro Leu Ile His Arg Gly	Tyr Tyr Val Arg	Ala

	50		55		60									
Arg	Ala	Val	Arg	His	Cys	Val	Arg	Ala	Phe	Leu	Glu	Gln	Ile	Gly
	65								70					75
Ala	Pro	Gln	Ala	Ala	Leu	Arg	Ala	Gln	Ile	Leu	Ser	Leu	Gly	Ala
	80								85					90
Gly	Phe	Asp	Ser	Leu	Tyr	Phe	Arg	Leu	Lys	Thr	Ala	Gly	Arg	Leu
	95								100					105
Ala	Arg	Ala	Ala	Val	Trp	Glu	Val	Asp	Phe	Pro	Asp	Val	Ala	Arg
	110								115					120
Arg	Lys	Ala	Glu	Arg	Ile	Gly	Glu	Thr	Pro	Glu	Leu	Cys	Ala	Leu
	125								130					135
Thr	Gly	Pro	Phe	Glu	Arg	Gly	Glu	Pro	Ala	Ser	Ala	Leu	Cys	Phe
	140								145					150
Glu	Ser	Ala	Asp	Tyr	Cys	Ile	Leu	Gly	Leu	Asp	Leu	Arg	Gln	Leu
	155								160					165
Gln	Arg	Val	Glu	Glu	Ala	Leu	Gly	Ala	Ala	Gly	Leu	Asp	Ala	Ala
	170								175					180
Ser	Pro	Thr	Leu	Leu	Leu	Ala	Glu	Ala	Val	Leu	Thr	Tyr	Leu	Glu
	185								190					195
Pro	Glu	Ser	Ala	Ala	Ala	Leu	Ile	Ala	Trp	Ala	Ala	Gln	Arg	Phe
	200								205					210
Pro	Asn	Ala	Leu	Phe	Val	Val	Tyr	Glu	Gln	Met	Arg	Pro	Gln	Asp
	215								220					225
Ala	Phe	Gly	Gln	Phe	Met	Leu	Gln	His	Phe	Arg	Gln	Leu	Asn	Ser
	230								235					240
Pro	Leu	His	Gly	Leu	Glu	Arg	Phe	Pro	Asp	Val	Glu	Ala	Gln	Arg
	245								250					255
Arg	Arg	Phe	Leu	Gln	Ala	Gly	Trp	Thr	Ala	Cys	Gly	Ala	Val	Asp
	260								265					270
Ile	Asn	Glu	Phe	Tyr	His	Cys	Phe	Leu	Pro	Ala	Glu	Glu	Arg	Arg
	275								280					285
Arg	Val	Glu	Asn	Ile	Glu	Pro	Phe	Asp	Glu	Phe	Glu	Glu	Trp	His
	290								295					300
Leu	Lys	Cys	Ala	His	Tyr	Phe	Ile	Leu	Ala	Ala	Ser	Arg	Gly	Asp
	305								310					315
Thr	Leu	Ser	His	Thr	Leu	Val	Phe	Pro	Ser	Ser	Glu	Ala	Phe	Pro
	320								325					330
Arg	Val	Asn	Pro	Ala	Ser	Pro	Ser	Gly	Val	Phe	Pro	Ala	Ser	Val
	335								340					345
Val	Ser	Ser	Glu	Gly	Gln	Val	Pro	Asn	Leu	Lys	Arg	Tyr	Gly	His
	350								355					360
Ala	Ser	Val	Phe	Leu	Ser	Pro	Asp	Val	Ile	Leu	Ser	Ala	Gly	Gly
	365								370					375
Phe	Gly	Glu	Gln	Glu	Gly	Arg	His	Cys	Arg	Val	Ser	Gln	Phe	His
	380								385					390
Leu	Leu	Ser	Arg	Asp	Cys	Asp	Ser	Glu	Trp	Lys	Gly	Ser	Gln	Ile
	395								400					405
Gly	Ser	Cys	Gly	Thr	Gly	Val	Gln	Trp	Asp	Gly	Arg	Leu	Tyr	His
	410								415					420
Thr	Met	Thr	Arg	Leu	Ser	Glu	Ser	Arg	Val	Leu	Val	Leu	Gly	Gly
	425								430					435
Arg	Leu	Ser	Pro	Val	Ser	Pro	Ala	Leu	Gly	Val	Leu	Gln	Leu	His
	440								445					450
Phe	Phe	Lys	Ser	Glu	Asp	Asn	Asn	Thr	Glu	Asp	Leu	Lys	Val	Thr
	455								460					465
Ile	Thr	Lys	Ala	Gly	Arg	Lys	Asp	Asp	Ser	Thr	Leu	Cys	Cys	Trp
	470								475					480
Arg	His	Ser	Thr	Thr	Glu	Val	Ser	Cys	Gln	Asn	Gln	Glu	Tyr	Leu
	485								490					495
Phe	Val	Tyr	Gly	Gly	Arg	Ser	Val	Val	Glu	Pro	Val	Leu	Ser	Asp
	500								505					510
Trp	His	Phe	Leu	His	Val	Gly	Thr	Met	Ala	Trp	Val	Arg	Ile	Pro
	515								520					525
Val	Glu	Gly	Glu	Val	Pro	Glu	Ala	Arg	His	Ser	His	Ser	Ala	Cys

	530		535		540
Thr Trp Gln Gly	Gly Ala Leu Ile Ala	Gly Gly Leu Gly Ala	Ser		
	545		550		555
Glu Glu Pro Leu	Asn Ser Val Leu Phe	Leu Arg Pro Ile Ser	Cys		
	560		565		570
Gly Phe Leu Trp	Glu Ser Val Asp Ile	Gln Pro Pro Ile Thr	Pro		
	575		580		585
Arg Tyr Ser His	Thr Ala His Val Leu	Asn Gly Lys Leu Leu	Leu		
	590		595		600
Val Gly Gly Ile	Trp Ile His Ser Ser	Ser Phe Pro Gly Val	Thr		
	605		610		615
Val Ile Asn Leu	Thr Thr Gly Leu Ser	Ser Glu Tyr Gln Ile	Asp		
	620		625		630
Thr Thr Tyr Val	Pro Trp Pro Leu Met	Leu His Asn His Thr	Ser		
	635		640		645
Ile Leu Leu Pro	Glu Glu Gln Gln Leu	Leu Leu Leu Gly Gly	Gly		
	650		655		660
Gly Asn Cys Phe	Ser Phe Gly Thr Tyr	Phe Asn Pro His Thr	Val		
	665		670		675
Thr Leu Asp Leu	Ser Ser Leu Ser Ala	Gly Gln			
	680		685		

&lt;210&gt; 10

&lt;211&gt; 519

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte Clone 1652936

&lt;400&gt; 10

Met Met Ser Lys	Asn Asp Gly Glu Ile Arg	Phe Gly Asn Pro Ala	
1	5	10	15
Glu Leu His Gly	Thr Lys Val Gln Ile Pro	Tyr Leu Thr Thr Glu	
	20	25	30
Lys Asn Ser Phe	Lys Arg Met Asp Asp	Glu Asp Lys Gln Glu Thr	
	35	40	45
Gln Ser Pro Thr	Met Ser Pro Leu Ala Ser	Pro Pro Ser Ser Pro	
	50	55	60
Pro His Tyr Gln	Arg Val Pro Leu Ser His	Gly Tyr Ser Lys Leu	
	65	70	75
Arg Ser Ser Ala	Glu Gln Met His Pro Ala	Pro Tyr Glu Ala Arg	
	80	85	90
Gln Pro Leu Val	Gln Pro Glu Gly Ser Ser	Ser Gly Gly Pro Gly	
	95	100	105
Thr Lys Pro Leu	Arg His Gln Ala Ser Leu	Ile Arg Ser Phe Ser	
	110	115	120
Val Glu Arg Glu	Leu Gln Asp Asn Ser Ser	Tyr Pro Asp Glu Pro	
	125	130	135
Trp Arg Ile Thr	Glu Glu Gln Arg Glu Tyr	Tyr Val Asn Gln Phe	
	140	145	150
Arg Ser Leu Gln	Pro Asp Pro Ser Ser Phe	Ile Ser Gly Ser Val	
	155	160	165
Ala Lys Asn Phe	Phe Thr Lys Ser Lys Leu	Ser Ile Pro Glu Leu	
	170	175	180
Ser Tyr Ile Trp	Glu Leu Ser Asp Ala Asp	Cys Asp Gly Ala Leu	
	185	190	195
Thr Leu Pro Glu	Phe Cys Ala Ala Phe His	Leu Ile Val Ala Arg	
	200	205	210
Lys Asn Gly Tyr	Pro Leu Pro Glu Gly Leu	Pro Pro Thr Leu Gln	

Pro Glu Tyr Leu	215	220	225
Gln Leu Phe Asp	230	235	240
Pro Arg Asp Leu	245	250	255
Asp Leu Pro Val	260	265	270
Ala Leu Lys Ser	275	280	285
Glu Asp Pro Ala	290	295	300
Pro Arg Ser Arg	305	310	315
Lys Arg Gly Glu	320	325	330
Thr His Ser Arg	335	340	345
Pro Ser Val Pro	350	355	360
Ala Leu Pro Pro	365	370	375
Glu Ala Glu Leu	380	385	390
Ala Glu Ser Ser	395	400	405
Pro Ser Lys Pro	410	415	420
Thr Glu Asn Gln	425	430	435
Ala Ala Thr Met	440	445	450
Lys Gln Lys Lys	455	460	465
Ala Asn Ala Val	470	475	480
Leu Lys Glu Val	485	490	495
Glu Gln Leu Arg	500	505	510
	515		

&lt;210&gt; 11

&lt;211&gt; 334

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte Clone 1710702

&lt;400&gt; 11

Met Phe Arg Trp	5	10	15
Ala Leu Cys Asn	20	25	30
Leu Thr Tyr Phe	35	40	45
Ser Ala Ala Pro	50	55	60
Ala Lys Glu Gly			

	65		70		75
His Trp Cys Val	Leu Pro Phe Arg Val	Leu Val Leu Thr Ser			
	80		85		90
His Arg Gly Ile	Gln Met Tyr Glu Ser	Asn Gly Tyr Thr Met Val			
	95		100		105
Tyr Trp His Ala	Leu Asp Ser Gly Asp	Ala Ser Pro Val Gln Ala			
	110		115		120
Val Phe Ala Arg	Gly Ile Ala Ala Ser	Gly His Phe Ile Cys Val			
	125		130		135
Gly Thr Trp Ser	Gly Arg Val Leu Val	Phe Asp Ile Pro Ala Lys			
	140		145		150
Gly Pro Asn Ile	Val Leu Ser Glu Glu	Leu Ala Gly His Gln Met			
	155		160		165
Pro Ile Thr Asp	Ile Ala Thr Glu Pro	Ala Gln Gly Gln Asp Cys			
	170		175		180
Val Ala Asp Met	Val Thr Ala Asp Asp	Ser Gly Leu Leu Cys Val			
	185		190		195
Trp Arg Ser Gly	Pro Glu Phe Thr Leu	Leu Thr Arg Ile Pro Gly			
	200		205		210
Phe Gly Val Pro	Cys Pro Ser Val Gln	Leu Trp Gln Gly Ile Ile			
	215		220		225
Ala Ala Gly Tyr	Gly Asn Gly Gln Val	His Leu Tyr Glu Ala Thr			
	230		235		240
Thr Gly Asn Leu	His Val Gln Ile Asn	Ala His Ala Arg Ala Ile			
	245		250		255
Cys Ala Leu Asp	Leu Ala Ser Glu Val	Gly Lys Leu Leu Ser Ala			
	260		265		270
Gly Glu Asp Thr	Phe Val His Ile Trp	Lys Leu Ser Arg Asn Pro			
	275		280		285
Glu Ser Gly Tyr	Ile Glu Val Glu His	Cys His Gly Glu Cys Val			
	290		295		300
Ala Asp Thr Gln	Leu Cys Gly Ala Arg	Phe Cys Asp Ser Ser Gly			
	305		310		315
Asn Ser Phe Ala	Val Thr Gly Tyr Asp	Leu Ala Glu Ile Arg Arg			
	320		325		330
Phe Ser Ser Val					

<210> 12  
 <211> 569  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte Clone 3239149

<400> 12	
Met Asp Pro Ala Glu	Ala Val Leu Gln Glu Lys Ala Leu Lys Phe
1	5 10 15
Met Asn Ser Ser Glu	Arg Glu Asp Cys Asn Asn Gly Glu Pro Pro
	20 25 30
Arg Lys Ile Ile Pro	Glu Lys Asn Ser Leu Arg Gln Thr Tyr Asn
	35 40 45
Ser Cys Ala Arg Leu	Cys Leu Asn Gln Glu Thr Val Cys Leu Ala
	50 55 60
Ser Thr Ala Met Lys	Thr Glu Asn Cys Val Ala Lys Thr Lys Leu
	65 70 75
Ala Asn Gly Thr Ser	Ser Met Ile Val Pro Lys Gln Arg Lys Leu
	80 85 90
Ser Ala Ser Tyr Glu	Lys Glu Lys Glu Leu Cys Val Lys Tyr Phe
	95 100 105

Glu	Gln	Trp	Ser	Glu	Ser	Asp	Gln	Val	Glu	Phe	Val	Glu	His	Leu
				110					115					120
Ile	Ser	Gln	Met	Cys	His	Tyr	Gln	His	Gly	His	Ile	Asn	Ser	Tyr
				125					130					135
Leu	Lys	Pro	Met	Leu	Gln	Arg	Asp	Phe	Ile	Thr	Ala	Leu	Pro	Ala
				140					145					150
Arg	Gly	Leu	Asp	His	Ile	Ala	Glu	Asn	Ile	Leu	Ser	Tyr	Leu	Asp
				155					160					165
Ala	Lys	Ser	Leu	Cys	Ala	Ala	Glu	Leu	Val	Cys	Lys	Glu	Trp	Tyr
				170					175					180
Arg	Val	Thr	Ser	Asp	Gly	Met	Leu	Trp	Lys	Lys	Leu	Ile	Glu	Arg
				185					190					195
Met	Val	Arg	Thr	Asp	Ser	Leu	Trp	Arg	Gly	Leu	Ala	Glu	Arg	Arg
				200					205					210
Gly	Trp	Gly	Gln	Tyr	Leu	Phe	Lys	Asn	Lys	Pro	Pro	Asp	Gly	Asn
				215					220					225
Ala	Pro	Pro	Asn	Ser	Phe	Tyr	Arg	Ala	Leu	Tyr	Pro	Lys	Ile	Ile
				230					235					240
Gln	Asp	Ile	Glu	Thr	Ile	Glu	Ser	Asn	Trp	Arg	Cys	Gly	Arg	His
				245					250					255
Ser	Leu	Gln	Arg	Ile	His	Cys	Arg	Ser	Glu	Thr	Ser	Lys	Gly	Val
				260					265					270
Tyr	Cys	Leu	Gln	Tyr	Asp	Asp	Gln	Lys	Ile	Val	Ser	Gly	Leu	Arg
				275					280					285
Asp	Asn	Thr	Ile	Lys	Ile	Trp	Asp	Lys	Asn	Thr	Leu	Glu	Cys	Lys
				290					295					300
Arg	Ile	Leu	Thr	Gly	His	Thr	Gly	Ser	Val	Leu	Cys	Leu	Gln	Tyr
				305					310					315
Asp	Glu	Arg	Val	Ile	Ile	Thr	Gly	Ser	Ser	Asp	Ser	Thr	Val	Arg
				320					325					330
Val	Trp	Asp	Val	Asn	Thr	Gly	Glu	Met	Leu	Asn	Thr	Leu	Ile	His
				335					340					345
His	Cys	Glu	Ala	Val	Leu	His	Leu	Arg	Phe	Asn	Asn	Gly	Met	Met
				350					355					360
Val	Thr	Cys	Ser	Lys	Asp	Arg	Ser	Ile	Ala	Val	Trp	Asp	Met	Ala
				365					370					375
Ser	Pro	Thr	Asp	Ile	Thr	Leu	Arg	Arg	Val	Leu	Val	Gly	His	Arg
				380					385					390
Ala	Ala	Val	Asn	Val	Val	Asp	Phe	Asp	Asp	Lys	Tyr	Ile	Val	Ser
				395					400					405
Ala	Ser	Gly	Asp	Arg	Thr	Ile	Lys	Val	Trp	Asn	Thr	Ser	Thr	Cys
				410					415					420
Glu	Phe	Val	Arg	Thr	Leu	Asn	Gly	His	Lys	Arg	Gly	Ile	Ala	Cys
				425					430					435
Leu	Gln	Tyr	Arg	Asp	Arg	Leu	Val	Val	Ser	Gly	Ser	Ser	Asp	Asn
				440					445					450
Thr	Ile	Arg	Leu	Trp	Asp	Ile	Glu	Cys	Gly	Ala	Cys	Leu	Arg	Val
				455					460					465
Leu	Glu	Gly	His	Glu	Glu	Leu	Val	Arg	Cys	Ile	Arg	Phe	Asp	Asn
				470					475					480
Lys	Arg	Ile	Val	Ser	Gly	Ala	Tyr	Asp	Gly	Lys	Ile	Lys	Val	Trp
				485					490					495
Asp	Leu	Val	Ala	Ala	Leu	Asp	Pro	Arg	Ala	Pro	Ala	Gly	Thr	Leu
				500					505					510
Cys	Leu	Arg	Thr	Leu	Val	Glu	His	Ser	Gly	Arg	Val	Phe	Arg	Leu
				515					520					525
Gln	Phe	Asp	Glu	Phe	Gln	Ile	Val	Ser	Ser	Ser	His	Asp	Asp	Thr
				530					535					540
Ile	Leu	Ile	Trp	Asp	Phe	Leu	Asn	Asp	Pro	Ala	Ala	Gln	Ala	Glu
				545					550					555
Pro	Pro	Arg	Ser	Pro	Ser	Arg	Thr	Tyr	Thr	Tyr	Ile	Ser	Arg	
				560					565					



<210> 13  
 <211> 123  
 <212> PRT  
 <213> Homo sapiens  
 <220>  
 <221> misc-feature  
 <223> Incyte Clone 3315936

<400> 13  
 Met Glu Phe Leu Glu Ile Gly Gly Ser Lys Pro Phe Arg Ser Tyr  
 1 5 10 15  
 Trp Glu Met Tyr Leu Ser Lys Gly Leu Leu Ile Phe Val Val  
 20 25 30  
 Asp Ser Ala Asp His Ser Arg Leu Pro Glu Ala Lys Lys Tyr Leu  
 35 40 45  
 His Gln Leu Ile Ala Ala Asn Pro Val Leu Pro Leu Val Val Phe  
 50 55 60  
 Ala Asn Lys Gln Asp Leu Glu Ala Ala Tyr His Ile Thr Asp Ile  
 65 70 75  
 His Glu Ala Leu Ala Leu Ser Glu Val Gly Asn Asp Arg Lys Met  
 80 85 90  
 Phe Leu Phe Gly Thr Tyr Leu Thr Lys Asn Gly Ser Glu Ile Pro  
 95 100 105  
 Ser Thr Met Gln Asp Ala Lys Asp Leu Ile Ala Gln Leu Ala Ala  
 110 115 120  
 Asp Val Gln

<210> 14  
 <211> 1957  
 <212> DNA  
 <213> Homo sapiens  
 <220>  
 <221> misc-feature  
 <223> Incyte Clone 016108

<400> 14  
 atttttgtca ctttctgtgt gaactaaagt gattcaatgt ctcttttga ttgcttctgt 60  
 acttcaagaa cacaagttga atcactcaga cctgaaaaac agtctgaaac cagtatccat 120  
 caatacttgg ttgatgagcc aaccctttcc tggtcacgtc catccactag agccagtga 180  
 gtactatggt ccaccaacgt ttctcactat gagctccaag tagaaatagg aagaggattt 240  
 gacaacttga cttctgtcca tcttgcacgg catactccca caggaacact ggtaactata 300  
 aaaattacaa atctggaaaa ctgcaatgaa gaacgcctga aagctttaca gaaagccgtg 360  
 attctatccc actttttccg gcatcccaat attacaactt attggacagt tttcactgtt 420  
 ggcagctggc tttgggttat ttctocattt atggcctatg gttcagcaag tcaactcttg 480  
 aggacctatt tccctgaagg aatgagtga actttaataa gaaacattct ctttggagcc 540  
 gtgagagggt tgaactatct gcacaaaaat ggctgtattc acaggagtat taaagccagc 600  
 catatcctca tttctgggtga tggcctagt accctctctg gcctgtccca tctgcatagt 660  
 ttggttaagc atggacagag gcatagggct gtgtatgatt tcccacagtt cagcacatca 720  
 gtgcagccgt ggttgagtcc agaactactg agacaggatt tacatgggtt atatgtgaag 780  
 tcagatatct acagtgttgg gatcacagca tgtgaattag ccagtgggca ggtgcctttc 840  
 caggacatgc atagaactca gatgctgtta cagaaactga aaggtcctcc ttatagccca 900  
 ttggatatca gtattttccc tcaatcagaa tccagaatga aaaattccca gtcagggtga 960  
 gactctggga ttggagaaa gttgcttgtc tccagtggaa ctcacacagt aaatagtgac 1020  
 cgattacaca caccatcctc aaaaactttc tctcctgcct tctttagctt ggtacagctc 1080  
 tgtttgcaac aagatcctga gaaaaggcca tcagcaagca gtttattgtc ccatgttttc 1140  
 ttcaaacaga tgaaagaaga aagccaggat tcaatacttt cactgttgcc tctgtcttat 1200  
 aacaagccat caatatcatt gcctccagt ttaccttgga ctgagccaga atgtgatttt 1260  
 cctgatgaaa aagactcata ctgggaattc tagggctgcc aaatcatttt atgtcctata 1320

```

tacttgacac tttctccttg ctgctttttc ttctgtattt ctaggtacaa ataccagaat 1380
tatacttgaa aatacagttg gtgcactgga gaatctatta tttaaaacca ctctgttcaa 1440
aggggacacca gttttagtgc cctctgtttc gcacagagta ctatgacaag gaaacatcag 1500
aattactaat ctagctagtg tcattttatc tggaattttt ttctaagctg tgactaactc 1560
tttttatctc tcaatataat ttttgagcca gtttaatttt ttcagtattt tgctgtccct 1620
tgggaatggg ccctcagagg acagtgtctc caagtacatc ttctcccaga ttctctggcc 1680
tttttaatga gctattgtta aaccaacagg ctagtttatc ttacatcaga cccttttctg 1740
gtagagggaa aatgtttgtg ctttcccttt ttcttctgtt aatacttatg gtaaacacta 1800
actgagcctc actcacatta aatgattcac ttgaaatata tacagaaatt gtaatttgct 1860
tttttttaaa aaagggggct aaagtaacac tttcctactt atgtaaatta tagatcctaa 1920
attcacgcac ccctgtggag ctcaataaag atttact 1957

```

&lt;210&gt; 15

&lt;211&gt; 2545

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte Clone 640521

&lt;400&gt; 15

```

gagccgagct gggggcgag agcgcgggag gcgggcgagg cgcggaacca gtcacccagg 60
ctgcagtgca gtggtgcgat ctcggtcag tcattgcaac cttcacctcc cggattcaag 120
tgattctcct gcctcagcct ccgagtagc tgggattaca ggtgcccacc accatgcca 180
ggtggctccg ctgcccgatg ggagtgcccc agtgtgtgtg atgaagctgg cgcagtcacc 240
atgtcatcat gtgtctctag ccagcccagc agcaaccggg ccgcccccca ggatgagctg 300
gggggagagg gcagcagcag cagcgaaagc cagaagccct gtgaggccct gcggggcctc 360
tcattcctga gcatccacct ggcatggag tccttcattg tggtcaccga gtgtgagccg 420
ggctgtgtgt tggacctcgg cttggcgagg gaccggcccc tggaggccga tggccaagag 480
gtcccccttg actcctccgg gtcccaggcc cgccccacc tctccggtcg caagtgtct 540
ctgcaagagc ggtcccaggg tgggctggca gccggtggca gcctggacat gaacggacgc 600
tgcatctgcc cgtcccctgc ctactcacc gtcagctccc cgcagtcctc gcctcggctg 660
ccccggcggc cgacagtggg gtctcaccac gtctccatca cgggtatgca ggactgtgtg 720
cagctgaatc agtataacct gaaggatgaa attggaagg gctcctatgg tgtcgtcaag 780
ttggcctaca atgaaaatga caatacctac tatgcaatga agtgtgtgtc caaaaagaag 840
ctgatccggc aggcgggctt tccacgtcgc cctccacccc gaggcacccg gccagctcct 900
ggaggctgca tccagcccag ggccccatt gagcaggtgt accaggaaat tgccatcctc 960
aagaagctgg accaccccaa tgtggtgaag ctggtggagg tcctggatga ccccaatgag 1020
gaccatctgt acatgggtgt cgaactggtc aaccaagggc ccgtgatgga agtgcccacc 1080
ctcaaaccac tctctgaaga ccaggcccgt ttctacttcc aggatctgat caaaggcatc 1140
gagtacttac actaccagaa gatcatccac cgtgacatca aaccttccaa cctcctgttc 1200
ggagaagatg ggcacatcaa gatcgctgac tttggtgtga gcaatgaatt caagggcagt 1260
gacgcgctcc tctccaacac cgtgggcacg ccgccttca tggcaccga gtcgctctct 1320
gagacccgca agatcttctc tgggaaggcc ttggatgttt gggctatggg tgtgacacta 1380
tactgctttg tctttggcca gtgcccattc atggacgagc ggatcatgtg ttacacagt 1440
aagatcaaga gtcaggccct ggaatttcca gaccagcccg acatagctga ggacttgaag 1500
gacctgatca cccgtatgct ggacaagaac cccgagtcga ggatcgtggt gccggaatc 1560
aagctgcacc cctgggtcac gaggcattgg gcggagccgt tgccgtcggg ggatgagaac 1620
tgacgcgtgg tcgaagtgaac tgaagaggag gtcgagaact cagtcaaaca cattcccagc 1680
ttggcaaccg tgatcctggg gaagaccatg atacgtaaac gtccttttgg gaacccattc 1740
gagggcagcc ggcgggagga acgctcactg tcagcgctg gaaacttgct cacgaagcaa 1800
ggcagcgaag acaacctcca gggcaccgac ccgcccccg tgggggagga ggaagtgtc 1860
ttgtgagagg cagtccctgc gtggaaagt gctgggcccc cgccccggc tccccgcac 1920
gcatgcatcc actgcgccg gaggagggca tggagcccga gtatgtgctt ggatcgctcg 1980
acctcgcatg cgcgcgcgt cgctctgtgg gggctgtgtc accgcgtttc catagcagca 2040
tgtcctacgg aaaccagca cgtgtgtaga gactcgtatg tcatctctgg ttattgttt 2100
tttctttgt gtgtttaaag gggacaaaaa aaaaaaaaaa aaggacttga ctccatgacg 2160
tcgaccgtgg ccgctggctg gctggacagg cgggtgtgag gagttgcaga cccaaacca 2220
cgtgcatttt gggacaattg ctttttaaaa cgtttttatg ccaaaaatcc ttcatgtgta 2280
ttttcagaac cagctcagat ataccaagt actgtgtgtg gggtttgaca actgtgaaa 2340
ggcgagcaga aaactccggc ggtctgaggc catggaggtg gttgctgcat ttgagaggga 2400
gtagggggct agatgtggct cctagtgcac accggaaacc atggcacctt ccagagccgt 2460

```

ggctctcaagg agtcagagca gggctggccc tcagtagctg cagggagctt tgatggcaac 2520  
 ttattttgtt aagaagggtt ttttt 2545

<210> 16  
 <211> 3034  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte Clone 1250171

<400> 16  
 tcctgagtct cgaggaggcc gcgggagccc gccggcggtg gcgcggcgga gacccggctg 60  
 gtataacaag aggattgcct gatccagcca agatgcagag cacttctaata catctgtggc 120  
 ttttatctga ttttttaggc caaggagcta ctgcaaacgt ctttcgtgga agacataaga 180  
 aaactggtga tttatttgct atcaaagtat ttaataacat aagcttcctt cgtccagtg 240  
 atgttcaaat gagagaattt gaagtgttga aaaaactcaa tcacaaaaat attgtcaaat 300  
 tatttgctat tgaagaggag acaacaacaa gacataaagt acttattatg gaattttgtc 360  
 catgtgggag tttatacact gttttagaag aaccttctaa tgcctatgga ctaccagaat 420  
 ctgaattctt aattgttttg cgagatgttg tgggtggaat gaatcatcta cgagagaatg 480  
 gtatagtgtca ccgtgatatac aagccaggaa atatcatgcy tgttataggg gaagatggac 540  
 agtctgtgtga caaactcaca gattttgttg cagctagaga attagaagat gatgagcagt 600  
 ttgtttctct gtatggcaca gaagaatatt tgcacctga tatgtatgag agagcagtg 660  
 taagaaaaga tcatcagaag aaatatggag caacagttga tctttggagc attggggtaa 720  
 cattttacca tgcagctact ggatcactgc catttagacc ctttgaaggg cctcgttaga 780  
 ataaagaagt gatgtataaa ataattacag gaaagccttc tgggtgcaata tctggagtac 840  
 agaaagcaga aatgggacca attgactgga gtggagacat gcctgtttct tgcagtcttt 900  
 ctggggtctc tcaggttcta cttaccctcg tcttgcaaaa catccttgaa gcagatcagg 960  
 aaaagtgttg ggttttgac cagttttttg cagaaactag tgatatactt caccgaatgg 1020  
 taattcatgt ttttcgcta caacaaatga cagctcataa gatttatata catagctata 1080  
 atactgctac tatatttcat gaactggtat ataaacaaac caaaattatt tcttcaaatc 1140  
 aagaacttat ctacgaaggg cgacgcttag tcttagaacc tggaggctg gcacaacatt 1200  
 tccctaaaac tactgaggaa aacctatat tttgtagtaag ccgggaacct ctgaatacca 1260  
 taggattaat atatgaaaaa atttccctcc cttaaagtaca tccacgttat gatttagacg 1320  
 gggatgctag catggctaag gcaataacag ggggtgtgtg tttatgcctgc agaattgcca 1380  
 gtaccttact gctttatcag gaattaatgc gaaaggggat acgatggctg attgaattaa 1440  
 ttaagatga ttacaatgaa actgttcaca aaaagacaga agttgtgatc acattggatt 1500  
 tctgtatcag aaacattgaa aaaactgtga aagtatatga aaagttagt aagatcaacc 1560  
 tggaaagcggc agagttaggt gaaatttcag acatacacac caaattgttg agactttcca 1620  
 gttctcaggg aacaatagaa accagtcttc aggatatcga cagcagatta tctccagggt 1680  
 gatcactggc agacgcatgg gcacatcaag aaggcactca tccgaaagac agaaatgtag 1740  
 aaaaactaca agtctgttga aattgcatga cagagattta ctatcagttc aaaaaagaca 1800  
 aagcagaacg tagattagct tataatgaag aacaaatcca caaatttgat aagcaaaaaac 1860  
 tgtattacca tgccacaaaa gctatgacgc actttacaga tgaatgtgtt aaaaagtatg 1920  
 aggcattttt gaataagtca gaagaatgga taagaaagat gcttcatctt aggaacagt 1980  
 tattatcgct gactaatcag tgttttgata ttgaagaaga agtatcaaaa tatcaagaat 2040  
 atactaatga gttacaagaa actctgcctc agaaaatgtt tacagcttcc agtggaaatca 2100  
 aacataccat gacccaatt tatccaagtt ctaacacatt agtagaaatg actcttggt 2160  
 tgaagaaatt aaaggaagag atggaagggg tgggttaaaga acttgctgaa aataaccaca 2220  
 ttttagaaag gtttggtctt ttaaccatgg atggtggcct tgcgaacgtt gactgtcttt 2280  
 agctttctaa tagaagttta agaaaagtgt ccgtttgcac aagaaaaataa cgcttgggca 2340  
 ttaaataaat gcttttatag atagtcactt gtttctacaa ttcagtattt gatgtggctg 2400  
 tgtaaatatg tacaatattg taaatacata aaaaatatac aaatttttg ctgctgtgaa 2460  
 gatgtaattt tatcttttaa catttataat tatatgagga aatttgacct cagtgtatcac 2520  
 gagaagaaag ccatgaccga ccaatatgtt gacatactga tctctactc tgagtggggc 2580  
 taataaagtt attttctctg accgcctact ggaataatatt ttaagtggaa ccaaaatagg 2640  
 catccttaca aatcaggaag actgacttga cagctttgtg aatggtagaa cgggtggctac 2700  
 tgtgagtgagg gagcagaacc gcaccactgt tatactggga taacaatttt tttgagaagg 2760  
 ataaagtgcc attattttat tttacaaggt gccagatcc cagttatcct tgtatccatg 2820  
 taatttcaga tgaattatta agcaaacatt ttaaagtga ttcattatta aaaactattc 2880  
 atttttttcc tttggccata aatgtgtaat tgtcattaaa attctaaggt cattttcaact 2940  
 gttttaagct gtatatcttct ttaattctgc ttactatttc atggaaaaaa ataaatttct 3000

caatttttaaa aaattttttt ataaaaaaaaa aaaa

3034

&lt;210&gt; 17

&lt;211&gt; 1337

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte Clone 1911587

&lt;400&gt; 17

```

gaaagctgtg ggaccatcct ggcaaccccc gtgtttggct gggttctagc gtaccgtctg 60
tgtggccggt gggggacctg cggtcggagt gggagggcca gtctgcaccc aagagggtga 120
agaggacggg ctttaggctg gaagcgctt agaggagcca tttttccagg tggggcccca 180
ggcagagggt cgcacaggga gcctggccat agtcgcgcac caggggagggt ggagcgctc 240
ccagacccga gccccgacc tcagccaaac ccattccttc tgccttgga ggccagagg 300
gactctgagc atcgaaagc aggatgcctg gtttgctttt atgtgaacca acagagcttt 360
acaacatcct gaatcaggcc acaaaactct ccagattaac agacccaac tatctctgtt 420
tattggatgt ccgttccaaa tgggagtatg acgaaagcca tgtgatcact gcccttcgag 480
tgaagaagaa aaataatgaa tatcttctcc cggagtctgt ggacctggag tgtgtgaagt 540
actgcgtggt gtatgataac aacagcagca ccctggagat actcttaaaa gatgatgat 600
atgattcaga ctctgatggt gatggcaaa atcttgtgcc tcaagcagcc attgagtatg 660
gcaggatcct gacccgcctc acccaccacc ccgtctacat cctgaaaggg ggctatgagc 720
gcttctcagg cacgtaccac tttctccgga ccagaagat catctggatg cctcaggaac 780
tggatgcatt tcagccatac cccattgaaa tcgtgccagg gaaggctctc gttggcaatt 840
tcagtcaagc ctgtgacccc aagattcaga aggacttgaa aatcaaagcc catgtcaatg 900
tctccatgga tacagggccc ttttttgca gcatgtctga caggcttctg cacatccgga 960
tagaagattc cccggaagcc cagattcttc ccttcttacg ccacatgtgt cacttcattg 1020
aaattcacca tcaccttggc tctgtcattc tgatcttttc caccaggggt atcagccgca 1080
gttggtccgc catcatagcc tacctcatgc atagtaacga gcagacctg cagaggctct 1140
gggcttatgt caagaagtgc aaaaacaaca tgtgtccaaa tcggggattg gtgagccagc 1200
tgctggaatg ggagaagact atccttgagg attccatcac aaacatcatg gatccgctct 1260
actgatcttc tccgaggccc accgaagggt actgaagagc ctcacctggg ggcattttgt 1320
gggtggaggg ccagagt 1337

```

&lt;210&gt; 18

&lt;211&gt; 1639

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte Clone 2079081

&lt;400&gt; 18

```

gacaaaagcc agacacattt caacatgagg gacccactga cagattgtcc gtataataaa 60
gtatacaaga acctaaagga gttttctcaa aatggagaga atttctgcaa acaggtcaca 120
tctgttcttc agcaaagggt aaacctggaa attagctatg ccaaaggact tcagaaactg 180
gcaagcaagc tgagcaaaagc attacagaac acgagaaaaa gttgtgttag cagtgcctgg 240
gcctgggcct cagagggaat gaaatccaca gcgacctgc atcaaaaact tggcaaagca 300
attgaattgg aagcaataaa accgacttat caagtcctaa atgtacaaga gaagaagaga 360
aaatcacttg acaatgaagt tgaaaagaca gcaaactctg tcattagcaa ctggaatcag 420
caaattaagg ccaagaagaa attaatggtt agtaccaga aacatgaagc acttttccag 480
cttgtagaaa gctccaagca atctatgact gagaaggaga agcgggaagc cctcaataaa 540
ctgacaaaat caactgaaaa gttgaaaag gaagatgaaa attactacca aaaaaacatg 600
gcgggttatt ctaccagact gaaatgggaa aacacactag agaactgcta ccagagcatt 660
ctggagctgg agaaggaaag aattcaactt ttatgcaata acttaaacca gtacagccaa 720
catatttctc tttttggcca aacctgacc acatgccaca cgcagattca ctgtgccatc 780
agcaagattg acattgaaaa agatatccag gctgtaattg aagaaactgc aattttatct 840
acagaaaaaa aatctgagtt cctgttaacg gattactttg aagaagatcc taacagtgca 900
atggataaag agagacgaaa gtctttacta aaaccaaaat tattgagact gcagagagac 960

```

```

attgaaaaag cctcaaaaaga caaggaaggc ctggaacgaa tgcttaaaac gtactccagc 1020
acctcctcct tctctgatgc aaagagccag aaagacacag cagcggttaat ggatgagaac 1080
aatttgaaac tagacctttt ggaagcgaac tcctacaaac tgtcatcaat gttagcagaa 1140
cttgagcaaa gacctcaacc cagccatcct tgtagtaatt ccatcttcag gtggagggaa 1200
aaggagcata ctcatagcta tgtgaaaata tctcggcctt ttttaaatgaa gagattagag 1260
aatattgtga gcaaggcatc ttctgggtggg cagagcaatc caggttcttc aactccagcc 1320
cctgggtgcag cccagctcag cagcagactt tgcaaggcct tgtattcttt tcaagccagg 1380
caagatgatg agttgaattt ggaaaagggt gacattgtga ttatacacga gaaaaaagaa 1440
gaaggatggt ggtttggatc tttgaatggg aaaaaaggcc attttcctgc cgcttatgtg 1500
gaggagttag cttcaaatgc tggcaacaca gctacaaagg cataaaacaa gactctgaac 1560
atactacctt cacactcggg aatcaacaat acagtgtggt tcaaataaga ataaagtgtc 1620
cttaccttta aaaaaaaaaa

```

<210> 19  
 <211> 1504  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte Clone 2472655

```

<400> 19
cgaaatcgta ggacttccga aagcagcggg ggcgtttgct tcaactgctt gaagtgtgag 60
tgccgcgaaga tgcgaaagggt ggttttgatc accggggcta gcagtggcat tggcctggcc 120
ctctgcaagc ggctgtctggc ggaagatgat gagcttcac tgtgtttggc gtgcaggaaac 180
atgagcaagg cagaagctgt ctgtgctgct ctgctggcct ctcacccac tgctgaggtc 240
acctattgtcc aggtggatgt cagcaacctg cagtcggtct tccgggcctc caaggaactt 300
aagcaaagggt ttcagagatt agactgtata tatctaaatg ctgggatcat gcctaatacca 360
caactaaata tcaaagcact tttctttggc ctcttttcaa gaaaagtgat tcatatgttc 420
tccacagctg aaggcctgct gaccaggggt gataagatca ctgctgatgg acttcaggag 480
tggtttgaga ccaatgtctt tggccatttt atcctgattc gggaactgga gcctctcctc 540
tgtcacagtg acaatccatc tcagctcatc tggacatcat ctgcagtgcc aaggaaatct 600
aatttcagcc tcgaggactt ccagcacagc aaaggcaagg aacctacag ctcttccaaa 660
tatgccactg accttttgag tgtggctttg aacaggaact tcaaccagca ggtctctat 720
tccaatgtgg cctgtccagg tacagcattg accaatttga catatggaat tctgcctccg 780
tttatatgga cgctgttgat gccggcaata ttgctacttc gcttttttgc aaatgcattc 840
actttgacac catataatgg aacagaagct ctggtatggc ttttccacca aaagcctgaa 900
tctctcaatc ctctgatcaa atatctgagt gccaccactg gctttggaag aaattacatt 960
atgacccaga agatggacct agatgaagac actgctgaaa aattttatca aaagtactg 1020
gaactggaaa agcacattag ggtcactatt caaaaaacag ataatcaggc caggctcagt 1080
ggctcatgcc tataattcca gcactttggg aggccaaagg agaaggatca cttgagacca 1140
ggagtccaag accagcctga gaaacatagt gagcccttgt ctctacaaaa agaaataaaa 1200
ataatagctg ggtgtggtgg catgcgcatg tagtcccagc tactcagaag gatgaggtgg 1260
gaggatctct tgaggctggg aggcagaggt tgcagtgagc tgagattgtg ccaactgcact 1320
ccagcctggg tgacagcgag accctgtctc aaaatatgta tatatttaat atatatataa 1380
aaccagagct gacaatgaca ctctggaaca ttgcatacct tctgtacatt ctgggggtaca 1440
tggatttcta ctgagttgga taatatgcat ttgtaataaa ctatgaacta tgaaaaaaa 1500
aaaa

```

<210> 20  
 <211> 3096  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte Clone 2948818

```

<400> 20
gggtgttctt ataacttaag ttcagttttt tctttcctgt gaggaagagg cagtttttta 60
aatgaagcca tctctgggga aatcgtattg attgtttag ctataacgg aatttttaaa 120

```

```

gtcttttagta tgttgaactg gaaatatag agatgcgttt gagatctact gtgagttgca 180
tcataaatatc aaaggactga agttataaaa gagaaaagag aagtttgctg ctaaaatgaa 240
tctgagcaat atggaatatt ttgtgccaca cacaaaaagg tactgaggat ttacccccca 300
aaaaaaattg tcaatgagaa ataaagctaa ctgatataca aaagcagagc ctgctctact 360
ggccatcatg cgtaaagggg tgctgaagga cccagagatt gccgatctat cctacaaaga 420
tgatcctgag gaacttttta ttggtttgca tgaaattggg catggaagtt ttggagcagt 480
ttattttgct acaaatgctc acaccagtga ggtggtggca attaagaaga tgcctatag 540
tggaagcag acccatgaga aatggcaaga tattcttaag gaagttaa atttacgaca 600
attgaagcat cctaatacta ttgagtacaa aggctgttac ttgaaagaac aactgcttg 660
gttggtgatg gaatattgct taggctcagc ctctgattta ttagaagttc ataaaaaacc 720
acttcaggaa gtggagatcg ctgccattac tcatggagcc ttgcatggac tagcctacct 780
acattctcat gcattgattc ataggatat taaagcagga aatattcttc taacagagcc 840
aggctcaggt aaactagctg attttggtc tgcttcaatg gcttctctcg ccaactcctt 900
cgtgggcaca ccttactgga tggctccaga ggtgatctta gctatggatg aaggacagta 960
tgatgggaaa gttgatattt ggtcacttgg catcacttgt attgaattgg cggaacggaa 1020
cccgccctt ttcaacatga atgcaatgag tgcccttatat cacattgccc agaagcagtc 1080
cccaacgtta cagtctaata aatggacaga ctcccttagg agatttgttg attactgctt 1140
gcagaaaaata cctcaggaaa ggccaacatc agcagaacta ttaaggcatg actttgttcg 1200
acgagaccgg ccactacgtg tctctattga cctcatagag aggacaaaag atgcagttcg 1260
tgagctagat aacctacagt accgaaaaat cttttccaag agacacggaa 1320
tggaaccttg aatgagtcac aggaggatga ggaagacagt gaacatggaa ccagcctgaa 1380
cagggaaatg gacagcctgg gcagcaacca ttccattcca agcatgtccg tgagcacagg 1440
cagccagagc agcagtgatg acagcatgca ggaagtcatg gacgagagca gttccgaact 1500
tgatcatgat cagcatgacg aaagcacaat caattccagc tctccgctcg tgcatagaa 1560
agatcatgta ttcataaggg atgagcgagg ccacggcgat cccaggcctg agcccgggcc 1620
taccagctca gttcagagcc aggccctcca ctaccggaac agagagcgct ttgccacgat 1680
caaatcagca tctttggtta cacgacagat ccatgagcat gagcaggaga acgagttcg 1740
ggaacagatg tcaggttata agcggatgag gcgccagcac cagaagcagc tgatcgccct 1800
ggagaacaag ctgaaggctg agatggacga gcaccgcctc aagctacaga aggaggtgga 1860
gacgcagtc aacaactcgt ccatcgagct ggagaagctg gccaaagaag aagtggctat 1920
catagaaaag gaggcaaaag tagctgcagc agatgagaag aagttccagc aacagatctt 1980
ggcccagcag aagaaagatt tgacaacttt cttagaaagt cagaagaagc agtataagat 2040
ttgtaaggaa aaaataaaaag aggaaatgaa tgaggaccat agcacacca agaaagagaa 2100
gcaagagcgg atctccaaac ataaagagaa cttgcagcac acacaggctg aagaggaagc 2160
ccacctctc actcaacaga gactgtacta cgacaaaaat tgctcgttct tcaagcggaa 2220
aataatgatc aagcggcacg aggtggagca gcagaacatt cgggaggaa taaataaaaa 2280
gaggaccag aaggagatgg agcatgccat gctaattccg cagcagagc ccaccggaga 2340
gctagagtac aggcagctgc acacgttaca gaagctacgc atggatctga tccgtttaca 2400
gcaccagacg gaactggaaa accagctgga gtacaataag aggcgagaaa gagaactgca 2460
cagaaagcat gtcattggaac ttcggcaaca gccaaaaaac ttaaaggcca tggaaatgca 2520
aattaaaaaa cagttttcag acacttgcaa agtacagacc aaacagtata aagcactcaa 2580
gaatcaccag ttggaagtta ctccaaagaa tgagcacaaa acaatcttaa agacactgaa 2640
agatgagcag acaagaaaac ttgccatttt ggcagagcag tatgaacaga gtataaatga 2700
aatgatggcc tctcaagcgt tacggctaga tgaggctcaa gaagcagaat gccaggcctt 2760
gaggctacag ctccagcagg aaatggagct ctcaacgcc taccagagca aaatcaagat 2820
gcaaacagag gcacaacatg aacgtgagct ccagaagcta gagcagagag tgtctctcg 2880
cagagcacac cttgagcaga agattgaaga ggagctggct gcccttcaga aggaacgcag 2940
cgagagaata aagaacctat tggaaaggca agagcgagag attgaaactt ttgacatgga 3000
gagcctcaga atgggatttg ggaatttggg tacattagat tttcctaagg aggactacag 3060
atgagattaa attttttgcc atttacaaaa aaaaaa 3096

```

&lt;210&gt; 21

&lt;211&gt; 1527

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte Clone 054191

&lt;400&gt; 21

```

ctccgcttga ggagaagcgc caagtgcgca tggggacgct atagcaattc gtttgctgtc 60
cttcctctcc ttcgaagatg acaaggccta ccatcgcttc ttctgcctt tgggccgtca 120

```

```

ggcagttggt tgggacccgc tccaaccctc ggttcttctt gcaatacagt ggatacaatt 180
tgtcatggct actctgagtg ttataggttc aagttcactt attgcctatg ctgtattcca 240
taataatacag aaatctccag agataagacc acttttttat ctgagcttct gtgacctgct 300
cctgggactt tgctggctca cggagacact tctctatgga gcttcagtag caaataagga 360
catcatctgc tataacctac aagcagttgg acagatattc tacatttctt catttctcta 420
caccgtcaat tacatctggt atttgtacac agagctgagg atgaaacaca ccagagtggt 480
acagagcaca tctccactgg tgatagatta tacttgtcga gttggtcaaa tggcctttgt 540
tttctcaagc ctgataacct tgctattgat gacacctgta ttctgtctgg gaaatactag 600
tgaatgtttc caaaacttca gtcagagcca caagtgtatc ttgatgcact caccaccatc 660
agccatggct gaacttccac cttctgccaa cacatctgtc tgtagcacac tttattttta 720
tggtatcgcc attttccctgg gcagctttgt actcagcctc cttaccatta tgggtcttact 780
tatccgagcc cagacattgt ataagaagtt tgtgaagtca actggctttc tggggagtga 840
acagtgggca gtgattcaca ttgtggacca acgggtgcgc ttctaccag tggccttctt 900
ttgctgctgg ggcccagctg tcattctaata gatcataaag ctgactaagc cacaggacac 960
caagcttcac atggcccttt atgttctcca ggctctaacg gcaacatctc aggggtctact 1020
caactgtgga gtatatggct ggacgcagca caaattccac caactaaagc aggaggctcg 1080
gcgtgatgca gatacccaga caccattatt atgctcacag aagagattct atagcagggg 1140
cttaaattca ctggaatcca cctgactttt tcctgccagt acttctacca ttttttgaaa 1200
ctacaatact ggaacatcca ggaactggag ttattctacg ctaatggatt ggaagaatg 1260
ttgggaaaag acatcttaaa tcttttctaa ctatgcccta aactgcagaa ctcaaaggaa 1320
atatagtgcc attgttagta gtcattctag atgaattggg agtatctctc cagttattcc 1380
cagattcact agtgatcctt aaagtctcta ttcagggaga ggaagacact ttccatctca 1440
gagatagact cgtgttacct tgatggatat tggatttgct taagtctctt ctagaaaaaa 1500
taaattctag attattaaaa aaaaaaa 1527

```

&lt;210&gt; 22

&lt;211&gt; 2948

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte Clone 1403604

&lt;400&gt; 22

```

aaagaaagggt cagccgcaag cgaacttagc actggctaca ccctcctcaa ttctggttgg 60
cgagatgcgc tcttcccggg agtgacgcac aagtgccggc ggaaggggaa gtccaggagc 120
atgggtggtt tttttccccc taccgaggtc cgtgagggtg gtgctaacca aggggaggct 180
cacaaccgtg acagactgcc attcctgagt ctcttctggc catgggcccc cggagccgtg 240
agcgtcgggc aggcgcggta cagaacacca acgacagcag cgccctcagc aagcgttccc 300
tggccgcgcg cgggtacgtg caggaccctt ttgcccgtt gctggttccg ggccgaggcg 360
gccgcgcacc gctcattcac cgaggctact acgtccgcgc acgcgcctg aggactgctg 420
tgcgcgcttt tttggagcag attggcgcgc cccaggccgc gcttcgcgcg cagatcttgt 480
ctctcggcgc tggcttcgac tcgctctatt ttcgcttaaa aaccgcgggc cgcctggccc 540
gggctgcagt ctgggaggtg gattttccgg acgtggcgcg gcgcaaagca gaaaggattg 600
gagagacgcc agagctgtgc gcgttaaccg ggcttttcca gaggggggag cccgcgtccg 660
cgctgtgctt tgagagcgca gactactgca tcctgggtct ggacttgccg cagctccagc 720
gagtggagga ggccctgggc gccgcggggc tcgacgcagc ctacccact ctgctcctgg 780
ccgagggcgt gctgacctac ctcgagccgg agagtgccgc ggccctcatc gcctgggcag 840
cccagcgttt tcctaatagcc cttttcgtgg tctatgagca gatgaggcct caagacgcct 900
ttggccagtt catgctgcaa cattttcggc agctaaactc cccctgcat ggccctggagc 960
gttttccctga cgtggaggcg cagcggcgcc gcttccctca agctggctgg accgcctgcg 1020
gtgccgtgga cataaatgaa ttctatcact gctttcttcc cgcagaagaa cgccggcggg 1080
tggaataat tgaacccttt gacgaatttg aggagtggca tctgaagtgc gccatttatt 1140
tcattctggc agcttctagg ggagacaccc tctccacac cctagtgtt ccatcctcag 1200
aggcatttcc tcgcgtaaat cctgcttcgc cttcaggggt attccttgc agcgtagtca 1260
gtagcgaggg ccaggtccca aacctgaaga gatatggcca cgctctgtc ttcttgagcc 1320
cagacgttat tctcagtga ggaggatttg gagagcagga ggggcggcac tgccgagtga 1380
gccagtttca cttgctctca agagattgtg actctgaatg gaaaggcagc caaataggca 1440
gttggtggag tggagttcag tgggatggac gcctttatca caccatgaca agactctcag 1500
agagtgggt tctggttctg ggaggagac tgtccccagt aagtccagcc ttgggggttc 1560
tccagcttca tttttttaag agtgaggata ataactga ggacctgaaa gtgacaataa 1620
caaaggcgtg ccgaaaggat gattccactt tgtgtgttg gcggcattca acaacagaag 1680

```

```

tgctcgtca gaatcaggaa tatttgtttg tgtatggggg tcgaagcgtg gtggaacctg 1740
tactaagtga ctggcatttc ctccatgtag ggacaatggc ttgggtcagg atcccagtg 1800
agggagaagt acctgaagcc cggcatttct acagtgcctg cacttggcaa gggggagccc 1860
ttattgctgg aggtctcggg gcttctgagg agccattgaa ctctgtgctc tttctgagac 1920
caatctcttg tggattcttc tgggagtcag tagacatcca gcctcccatt accccaaggt 1980
actcccacac agctcatgtg ctcaatggaa agctgttact gggtggaggg atctggattc 2040
attcctcctc ctaattattca gtgactgtga tcaatttgac tacaggattg agctctgagt 2100
atcagattga cacaacatat gtgccatggc cattaatgtt acacaacat actagtatcc 2160
ttcttcctga agagcaacag ctccgtgctc ttggaggtgg tgggaactgc ttttcctttg 2220
gtacctactt caacccccat acagtcacat tagaccttc ttccttaagt gctgggcagt 2280
aaggactgga ctaattattca ggaccacta aagtagacaa taaagttttc cacaatatag 2340
atgacctct agctatagat actgccactc ctcccttccc catccttttt ttccttagc 2400
actattcagt gcaaaaagtg aaaaagggtg gtaaaatagg taaaatacct agaacaatc 2460
actacagaaa acagctgaag acagtggcca tgcagtcoga gaggagtagt ggtctgcctc 2520
taattttcta atctaagttc gtttattgag ttacagtggg ctttagtaaa gtaaaacaat 2580
ttcccaatcc caggccttgt gatttgagat ggtacctag aaaaagtta acgcagttcc 2640
gtggttgaat atatttgaga tggtaacctt gaaaaagttt cacgcagatc cttggttgaa 2700
tatagttgag ggagcgtagt attgacaatt ctcatgtag gaaacctgaa atgaacacag 2760
tcacagtttg attaaaacat tgcctgtttt gtgcaacag aaaactcoga tagttttaac 2820
aacaggaaac actttagga cttcctttac caacatactt tttaaatgtt ttgctattgg 2880
ttccatattt atttagattt ataagtgtca ataaagcaaa cttttgatgc ctcaaaaaaa 2940
aaaaaaa

```

<210> 23  
 <211> 1808  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte Clone 1652936

```

<400> 23
gagagtatta aatgtgaatt gcctctgcct cgctttatga tgtcaaagaa tgatggtgag 60
atacagatttg ggaacccagc tgagctgcat ggaactaagg ttcagattcc atatttaact 120
acagaaaaaa attccttcaa aagaatggac gatgaggata aacaggaaac acagtctccc 180
acgatgtcac ccctcgcctc ccctccttct tcccgcctc attaccagag ggtgcccttg 240
agccatggct acagcaaaact gcggagcagc gcagaacaga tgcattccagc accttatgaa 300
gctaggcagc cccttgtcca gcccagggga tctcatcag ggggcccagg aaccaagccc 360
cttcggcatc aggccttccct tatccggtcc ttttcagtgg agagggaaact acaggataac 420
agcagttacc ccgacgaacc ctggaggata acagaagaac agcgcgagta ctatgtcaat 480
cagttcogat cccttcagcc agacccaagc tctttcattt cagggttctgt ggccaagaac 540
ttcttcacca aatcaaagct ttccattcca gaactctcct atatatggga gcttagtgat 600
gctgactgtg atggagccct gacctgcct gagttctgtg ctgctgttca tctcattgtg 660
gctcggaaga acggctaccc attgcctgag ggccctccct caactctgca gccagaatac 720
ctgcaggcag cttttcctaa gcccaaatgg gactgtcaat tatttgattc ttattctgag 780
tcactgccgg caaatcaaca acctcgtgac ttgaatcgga tggagacatc tgttaaagac 840
atggctgacc ttctgttccc taaccaggat gtaactagtg atgacaaaca agctttgaaa 900
agtactatca atgaagcctt accaaaggac gtgtctgagg atccagcaac tcccaaggat 960
tccaacagtc tcaaagcaag accaagatcc agatcttact ctagcacctc catagaagag 1020
gccatgaaaa ggggcgagga ccctcccacc ccgccacctc ggccacagaa aacccattcc 1080
agagcctcct ccttgatctt gaataaagtc ttccagccca gtgtgccagc taccagtgca 1140
ggattgttac cccaccacc tgcgctccct ccaagacctt gtccatcaca gtctgaacaa 1200
gtgtcggagg ccgagttact cccacagctg agcagagccc catcccaggc tgcagaaaag 1260
agtccagcaa agaaggatgt actgtattct cagccacctt caaagcccat tcgtaggaaa 1320
ttcagaccag aaaaccaagc tacagaaaac caagagcctt ccaactgctgc aagtgggcca 1380
gcttctgcgg caaccatgaa accgcatcca acagtccaaa agcagttctc caaacagaag 1440
aaggccattc aaactgctat ccgcaaaaat aaagaggcaa acgcagtgct ggctcggctg 1500
aacagtgagc tccagcagca gctcaaggag gtcatcaag aacgaattgc attggaacaa 1560
caattggaac aacttcgtcc ggtcactgtg ttgtgacccc ccatgggttc aagtgcagat 1620
gggtgacctt gctgtccaa gctgtgagca gcaaaatata atccatatga ctttttctct 1680
gatgtttgac tgtgtcaaaa gctgtgagca gcaaaatata atccatatga ctttttctct 1740
tgtatagact taaaaaaa aaaatagatc ttaattaag cggtcgcaag cttattccct 1800

```